arXiv:hep-th/0406072v1 8 Jun 2004

Abelian and non-Abelian branes in WZW models and gerbes

Krzysztof Gawędzki*

Laboratoire de Physique, ENS-Lyon,
46, Allée d'Italie, F-69364 Lyon, France

Abstract

We discuss how gerbes may be used to set up a consistent Lagrangian approach to the WZW models with boundary. The approach permits to study in detail possible boundary conditions that restrict the values of the fields on the worldsheet boundary to brane submanifolds in the target group. Such submanifolds are equipped with an additional geometric structure that is summarized in the notion of a gerbe module and includes a twisted Chan-Paton gauge field. Using the geometric approach, we present a complete classification of the branes that conserve the diagonal current-algebra symmetry in the WZW models with simple, compact but not necessarily simply connected target groups. Such symmetric branes are supported by a discrete series of conjugacy classes in the target group and may carry Abelian or non-Abelian twisted gauge fields. The latter situation occurs for the conjugacy classes with fundamental group $\mathbb{Z}_2\times\mathbb{Z}_2$ in $SO(4n)/\mathbb{Z}_2$. The branes supported by such conjugacy classes have to be equipped with a projectively flat twisted $U(2)$ gauge field in one of the two possible WZW models differing by discrete torsion. We show how the geometric description of branes leads to explicit formulae for the boundary partition functions and boundary operator product coefficients in the WZW models with non-simply connected target groups.

1 Introduction

Boundary conformal field theories play an important role in the study of two- or $(1+1)$-dimensional critical phenomena in finite geometries and in the understanding of branes in string theory. In the latter context much of the original work has been done in the Lagrangian approach where one considers tense strings moving in flat or nearly flat space-times, with ends restricted to special submanifolds called Dirichlet branes or D-branes [61]. From the worldsheet point of view, this amounts to the study of conformally invariant boundary sigma models with flat or nearly flat target spaces and with fields on the worldsheet boundary restricted to take values in the brane submanifolds of the target. The latter are chosen in a way that assures that the boundary theory still has (half of) the conformal symmetry. Such boundary sigma models constitute, at least perturbatively, examples of boundary conformal field theories. On the non-perturbative level, the field theories of this type may be analyzed in algebraic terms with the powerful 2d conformal symmetry toolkit [20], as initiated in [16]. In this way, boundary conformal field theory provides a formulation of the stringy or D-geometry [22]. Understanding how this geometry modifies the standard geometry requires also a translation of as many of its aspects as possible to a more standard Lagrangian geometric language. This program has been implemented to a certain degree for the Calabi-Yau sigma models and the $N = 2$ supersymmetric boundary conformal field theories, see [22, 23, 5], where, due to supersymmetry, the relation between the stringy and the classical geometry may be traced a long way and is reflected in the mirror symmetry [48].

---

*membre du C.N.R.S.

Similar question may be studied for the Wess-Zumino-Witten (WZW) model, a conformally invariant sigma model with a group manifold as the target [71]. Here, it is the rich current-algebra symmetry that allows to trace the relation between the stringy and the classical geometry. In the WZW model, one may access the stringy geometry via the exact solution of the corresponding boundary conformal field theory. On the other hand, one may interpret this solution within the Lagrangian approach. In particular, the WZW theories provide an important laboratory for studying stringy geometry of curved D-branes with non-trivial fluxes of the Kalb-Ramond B-field [67]. In the simplest case, the so called (maximally) symmetric D-branes in the WZW models with compact targets correspond to a discrete series of the integrable conjugacy classes in the group manifold [47, 2]. These are the conjugacy classes that contain elements $\mathrm{e}^{2\pi\mathrm{i}\lambda/\mathsf{k}}$ where k is the level of the model (related to its coupling constant) and λ is the highest weight of a chiral current-algebra primary field of the theory. For the so called diagonal WZW models with simply connected target groups, there is a unique integrable conjugacy class for each λ and a unique maximally symmetric conformal boundary condition associated to it [2], see also [37]. This results in a one-to-one correspondence between the primary fields and the boundary conditions, as predicted in [16]. The situation is more complex for the non-diagonal WZW models corresponding to non-simply connected target groups. The set of maximally symmetric conformal boundary conditions and the exact solution for the boundary model have been described in [30, 32, 70] in terms of the so called simple current structure [50], see also [7, 59, 34]. This description exposes a more complicated relation between the integrable conjugacy classes and the boundary conditions, with occurrence of non-trivial multiplicities of the latter for a fixed conjugacy class and with a possibility of spontaneous generation of non-Abelian gauge-type structure [31]. In particular, ref. [32]stressed the role of the finite group cohomology in the boundary model. The aim of the present paper is to provide a geometric picture of the classification of the symmetric branes for the WZW models with non-simply connected targets by carefully setting up the Lagrangian approach to such models in the presence of boundaries.

The characteristic feature of the WZW model in the Lagrangian formulation is the presence of the coupling to a topologically non-trivial Kalb-Ramond background 2-form field B on the target group. This field is defined locally, only with its exterior derivative $H = dB$, equal to the standard invariant 3-form on the group, making sense globally. As noted in [71], the fact that the 3-form H is not exact so that there is no global B

field allows a consistent definition of the classical amplitudes in the bulk theory (i.e. on worldsheets without boundary) only if the periods of the 3-form H are in $2\pi\mathbb{Z}$. This leads to the quantization of the the coupling constant of the WZW model that was analyzed in [71] for simply connected target groups and in [27] for non-simply connected ones. The careful Lagrangian formulation of the theory permitted a direct calculation of the quantum spectra of the bulk WZW models [42, 27].

An idea of how to define the bulk amplitudes for a general sigma model coupled to a closed 3-form H on the target with periods in $2\pi\mathbb{Z}$ was proposed in [3]. This proposal was reformulated in terms of the 3^{rd} degree Deligne cohomology in [36]. The recent years have brought a realization that such cohomology classifies geometric objects over the target manifolds called (Abelian) bundle gerbes with connection [56, 19, 44][1]. The closed 3-form H provides the curvature form of the gerbe. In the presence of boundaries, the B-field ambiguities lead to new phenomena and result in the quantization of possible D-brane boundary conditions first observed in [47] and [2] for the case of the WZW models with simply connected targets. The geometric language of gerbes appears quite useful in dealing with the intricacies of the open string amplitudes in the presence of topologically non-trivial B-fields. In particular, the Lagrangian description of the Chan-Paton coupling of the ends of the open string to a non-Abelian Yang-Mills field on the D-brane leads in a natural way to the notion of gerbe modules, as first noted in ref. [46] inspired by [29], see also [17] (in [46] such modules were viewed as modules over the Azumaya algebras). The gerbe modules also appear in a definition [10] of the twisted K-theory groups containing the Ramond-Ramond charges of the supersymmetric D-branes [72, 54].

The construction of the bundle gerbes with curvature proportional to the invariant 3-form over the $SU(2)$ group may be traced back to [36] and predates the geometric definition of the notion. The bundle gerbes over $SU(N)$ were constructed in [19] and, independently but later, in [39]. The extension of this construction to other simple, compact, simply connected groups required overcoming additional difficulties and was achieved in [53], see [15, 8] for a related work. The bundle gerbes over non-simply connected groups were described in [36] for the $SO(3)$ group, in [39] for the groups covered by $SU(N)$ and in [40] for all non-simply connected, simple, compact Lie groups. As is well known, see e.g. [49], pushing down a line bundle from manifold M to its quotient by free action of a finite group Z requires solving a cohomological equation $\delta W = V$ that expresses a 2-cocycle V on group Z as a boundary, with the cohomology class of V representing the obstruction. Similar considerations for gerbes [68, 62] lead to a cohomological equation $\delta V = U$ in one degree higher, with the obstruction to pushing down a gerbe to the quotient space represented by the cohomology class of the 3-cycle U on Z. Calculating the 3-cocycles U and solving the equation $\delta V = U$ whenever possible for all quotients G' of simply connected, simple, compact groups G by subgroups Z of their center was the essence of the work done in [40]. We shall crucially depend in the present paper on those results. In [39], the gerbes over $SU(N)$ and its quotients were used to classify the symmetric D-branes of the WZW models with these groups as the target. Such branes consisted of an integrable conjugacy class in the group equipped with additional geometric structure which included an Abelian gauge field on the conjugacy class. Pushing down the line bundle carrying the gauge field from a conjugacy class in $SU(N)$ to a conjugacy class in the quotient group required solving a cohomological equation $\delta W = V$, as discussed above.

---

[1] see [43] and [14] for the categorial avatars of bundle gerbes

Below, we shall extend the geometric classification of the symmetric D-branes to WZW models with target groups that are simple, compact but not necessarily simply connected. As in the $SU(N)$ case, such branes are supported by integrable conjugacy classes. The general case will require to include in the additional structure a projectively flat twisted non-Abelian gauge field on the conjugacy class. The adequate formulation is based on the notion of gerbe modules, also called twisted vector bundles [46, 51, 10]. More exactly, we shall classify, up to isomorphism, the possible gerbe modules over the integrable conjugacy classes $\mathcal{C}$ with the gerbe over $\mathcal{C}$ obtained from the gerbe over the group by restriction. For simply connected groups there is a unique 1-dimensional gerbe module corresponding to a symmetric brane over each integrable conjugacy class. All higher dimensional ones are its direct sums. The construction of the gerbe modules over integrable conjugacy classes $\mathcal{C}'$ in the non-simply connected quotient groups $G' = G/Z$ leads to a cohomological equation $\delta\mathcal{W} = \mathcal{V}$ expressing a 2-cocycle $\mathcal{V}$ on the fundamental group of the conjugacy class $\mathcal{C}'$ as a boundary, similarly as for pushing to quotients usual line bundles. The cohomology class of $\mathcal{V}$ is the obstruction to the existence of a 1-dimensional gerbe module over $\mathcal{C}'$ giving rise to a symmetric brane. If the obstruction class is trivial then such 1-dimensional modules exist and different ones differ by tensor multiplication by flat line bundles over $\mathcal{C}'$. Higher dimensional modules are then direct sums of 1-dimensional ones. If the obstruction class is non-trivial, there are still matrix-valued solutions of the equation $\delta\mathcal{W} = \mathcal{V}$ and they give rise to higher dimensional gerbe modules over the conjugacy classes $\mathcal{C}'$ that do not decompose into a direct sum of 1-dimensional ones. This mimics the situation for line bundles which may always be pushed down to vector bundles over regular discrete quotients. The matrix-valued solutions $\mathcal{W}$ give rise to symmetric branes carrying a projectively flat twisted non-Abelian gauge field. Such a situation occurs for the conjugacy classes with fundamental group $\mathbb{Z}_2 \times \mathbb{Z}_2$ in the $SO(4n)/\mathbb{Z}_2$ groups. These are groups that admit two non-equivalent gerbes distinguished by discrete torsion [69, 68], as first realized in [27] where the ambiguity was identified as periodic vacua. For one of those gerbes, the possible gerbe modules leading to symmetric branes are direct sums of 2-dimensional modules that cannot be decomposed further. Similar spontaneous enhancement of the gauge symmetry has been previously observed at orbifold fixed points of the Calabi-Yau sigma models in [21, 24]. Within the algebraic approach based on simple current structure to non-diagonal boundary WZW and coset models, it was related in [31] to the presence of projective representations of the simple current symmetries. We recover such representations in the description of the 2-dimensional gerbe modules corresponding to the D-branes in $SO(4n)/\mathbb{Z}_2$.

The geometric constructions, although based on classical considerations, lead through geometric quantization to simple expressions for the boundary partition functions and the boundary operator product coefficients in the quantum WZW models with non-simply connected target groups, as noted in [39]. Such expressions result from a realization of the open string states of the model with a non-simply connected target group $G' = G/Z$ as the states of the model with the covering group G target that are invariant under an action of the fundamental group Z. The existence of such an action and the above picture become clear in the geometric realization of the states of the models, see

Sect. 9.2 of [39]. Although they may be, and will be, formulated within a more standard algebraic approach, they employ the cohomological structures that appear naturally in the geometric classification of the branes discussed here showing the pertinence of the latter for the quantum theory.

The paper is organized as follows. In Sect. 2, we present the notions of bundle gerbes and gerbe modules in a relatively pedestrian fashion, with stress on the local description. How those notions are employed to define the contributions to the action functional of the topologically non-trivial B field and the Chan-Paton coupling to the non-Abelian gauge fields on the branes in the presence of such a B field is the subject of Sect. 3. The next two sections are the most technical ones. In Sect. 4 we recall briefly the construction of the gerbes over Lie groups relevant for the treatment of the B field contributions in the WZW models. We discuss the case of simple compact groups, both simply connected [53] and non-simply connected [40]. The symmetric branes and their geometric classification are studied in Sect. 5, first for the simply connected groups (in Sect. 5.1) and then for the non-simply connected ones (in Sect. 5.2). The latter discussion, culminating in considerations leading to the cohomological equation $\delta\mathcal{W}=\mathcal{V}$ constitutes the main part of the present work. We have not found yet a more concise way to describe the solutions $\mathcal{W}$ for all cases with trivial cohomology class of $\mathcal{V}$ and have opted for tabulating the results in Appendix A where the complete list of symmetric branes carrying Abelian gauge fields is given. In Sect. 6, we discuss the special case of symmetric branes in $SO(4n)/\mathbb{Z}_2$ that carry a twisted non-Abelian gauge field. Sect. 7 contains a discussion of the boundary space of states in the WZW models, of the boundary partition functions and the boundary operator product. It completes the discussion of Sects. 9 and 10 of [39]. Conclusions summarize the results of the paper and draw perspectives for the future research. Appendix B provides a direct check that the action of elements of the group center on the multiplicity spaces of the boundary theory with simply connected target is well defined and Appendix C shows how the general formulae of Sect. 7 permit to obtain the boundary partition functions and the boundary operator product coefficients for the WZW model with the $SO(3)$ target.

Acknowledgements. This work has stemmed from research conducted initially in collaboration with Nuno Reis. It was partially done in framework of the European contract EUCLID/HPRN-CT-2002-00325.

2 Abelian bundle gerbes and gerbe modules

2.1 Local data

We start by a quick introduction to Abelian bundle gerbes and gerbe modules. It is simplest to specify such objects by local data. For the bundle gerbes, the local approach was developed long before the geometric approach of [56, 57], see [3, 36]. For bundle-gerbe modules [10], also called Azumaya algebra modules or twisted vector bundles, the local approach was discussed in [46], see also [51].

Let (O_i) be a good open covering of a manifold M (i.e. such that O_i and all non-empty intersections $O_{i_1\dots i_n}=O_{i_1}\cap\dots\cap O_{i_n}$ are contractible). We shall call a family (B_i,A_{ij},g_{ijk}) local data for an Abelian hermitian bundle gerbe with connection (in short, **gerbe local data**) if B_i are 2-forms on O_i, $A_{ij} = -A_{ji}$ are 1-forms on O_{ij} and $g_{ijk} = g^{sgn(\sigma)}_{\sigma(i)\sigma(j)\sigma(k)}$ are $U(1)$ valued functions on O_{ijk} such that

$$\begin{aligned} B_j - B_i \ &= \ dA_{ij} && \text{on } O_{ij}\,, && (2.1,a)\\ A_{jk} - A_{ik} + A_{ij} \ &= \ i\, g_{ijk}^{-1} dg_{ijk} && \text{on } O_{ijk}\,, && (2.1,b)\\ g_{jkl}\, g_{ikl}^{-1}\, g_{ijl}\, g_{ijk}^{-1} \ &= \ 1 && \text{on } O_{ijkl}\,. && (2.1,c) \end{aligned}$$

The global closed 3-form H equal to dB_i on O_i is called the **curvature** of the corresponding data. Local data with curvature H exist if and only if the 3-periods of H are in $2\pi\mathbb{Z}$. Writing $g_{ijk} = \mathrm{e}^{2\pi \mathrm{i} f_{ijk}}$ for real-valued functions f_{ijk} one obtains an integer-valued 3-cocycle

$$n_{ijkl} \ = \ f_{jkl} - f_{ikl} + f_{ijl} - f_{ijk} \qquad \text{on } O_{ijkl}$$

that defines a cohomology class in $H^3(M,\mathbb{Z})$, called the **Dixmier-Douady** (DD) **class** of the gerbe local data. The image of the DD-class in $H^3(M,\mathbb{R})$ coincides with the de Rham cohomology class of H.

One says that two families (B_i, A_{ij}, g_{ijk}) and $(B'_i, A'_{ij}, g'_{ijk})$ of gerbe local data are **equivalent** if

$$\begin{aligned} B'_i \ &= \ B_i + d\pi_i\,, && (2.2,a)\\ A'_{ij} \ &= \ A_{ij} + \pi_j - \pi_i - i\,\chi_{ij}^{-1} d\chi_{ij}\,, && (2.2,b)\\ g'_{ijk} \ &= \ g_{ijk}\,\chi_{ik}\,\chi_{jk}^{-1}\,\chi_{ij}^{-1}\,, && (2.2,c) \end{aligned}$$

where π_i are 1-forms on O_i and $\chi_{ij} = \chi_{ji}^{-1}$ are $U(1)$-valued functions on O_{ij}. The DD-classes of equivalent local data coincide but one may have non-equivalent local data that correspond to the same DD-class.

Given (B_i, A_{ij}, g_{ijk}) as above, we shall call a family (Π_i, G_{ij}) local data for an N-dimensional hermitian gerbe module with connection (in short: **gerbe-module local data**) if Π_i are $u(N)$-valued[2] 1-forms on O_i and G_{ij} are $U(N)$-valued maps on O_{ij} such that

$$\begin{aligned} \Pi_j - G_{ij}^{-1}\Pi_i\, G_{ij} \ - i\, G_{ij}^{-1} dG_{ij} + A_{ij} \ &= \ 0 && \text{on } O_{ij}\,, && (2.3,a)\\ G_{ik}\, G_{jk}^{-1}\, G_{ij}^{-1}\, g_{ijk} \ &= \ 1 && \text{on } O_{ijk}\,. && (2.3,b) \end{aligned}$$

Recall that local data of a hermitian vector bundle with connection satisfy similar relations with $A_{ij} = 0$ and $g_{ijk} = 1$. Two families of gerbe-module local data (Π_i, G_{ij}) and (Π'_i, G'_{ij}) are said to be **equivalent** if

$$\begin{aligned} \Pi'_i \ &= \ H_i^{-1}\,\Pi_i\, H_i \ + i\, H_i^{-1} dH_i\,, && (2.4,a)\\ G'_{ij} \ &= \ H_i^{-1}\, G_{ij}\, H_j && (2.4,b) \end{aligned}$$

for $U(N)$-valued functions H_i on O_i. Given gerbe-module local data (Π_i, G_{ij}) relative to gerbe local data (B_i, A_{ij}, g_{ijk}) and equivalent local gerbe data $(B'_i, A'_{ij}, g'_{ijk})$, the family $(\Pi_i - \pi_i,\ G_{ij}\chi_{ij}^{-1})$ provides a gerbe-module local data relative to $(B'_i, A'_{ij}, g'_{ijk})$. We shall call such data **induced** from (Π_i, G_{ij}) by the equivalence of local gerbe data.

---

[2] $u(N)$ denotes the Lie algebra of hermitian $N \times N$ matrices

Gerbe-module local data exist only if and only if the DD-class of the associated gerbe local data is pure torsion, i.e. if the curvature H is an exact form, a very stringent requirement. The only if part may be be easily deduced from (2.3) by taking its trace and exterior derivative to infer that $B_i + \frac{1}{N}\,\mathrm{tr}\,\Pi_i$ define a global form on M whose exterior derivative is equal to H. The if part may be inferred from the discussion in [10].

It is sometimes convenient to reduce the structure group of a gerbe module (as for the standard vector bundles) from $U(N)$ to a compact group G assumed here to contain a distinguished central $U(1)$ subgroup. This is done by requiring in the definitions above that Π_i are $\mathbf{g}$-valued forms, where $\mathbf{g}$ is the Lie algebra of G, and G_{ij} and H_i are G-valued. One may then return to the previous situation by composition with an N-dimensional unitary representation U of G acting as identity on the $U(1)$ subgroup of G.

2.2 Geometric definitions

One may construct geometric objects from local data (B_i, A_{ij}, g_{ijk}). Let $Y = \sqcup_i O_i$ be the disjoint union of O_i and π be the natural map $(m,i) \mapsto m$ from Y to M. Local 2-forms B_i define a global 2-form B on Y such that

$$dB \;=\; \pi^* H\,. \tag{2.5}$$

Denote by $Y^{[n]}$ the n-fold fiber product $Y \times_M \cdots \times_M Y$. Clearly, $Y^{[n]} = \underset{(i_1,\dots,i_n)}{\sqcup} O_{i_1\dots i_n}$ and we shall use the notation $(m, i_1, \dots, i_n)$ for its elements. Consider the trivial line bundle $L = Y^{[2]} \times \mathbb{C}$ over $Y^{[2]}$ with the hermitian structure inherited from $\mathbb{C}$. It may be equipped with a (unitary) connection with local connection forms equal to A_{ij} on O_{ij}. The curvature form F of this connection satisfies

$$F \;=\; p_2^* B \,-\, p_1^* B \tag{2.6}$$

where p_i denote the projections in $Y^{[n]}$ on the $i^{\,\mathrm{th}}$ factor. Finally, L may be equipped with a groupoid multiplication

$$L_{(y,y')} \times L_{(y',y'')} \;\xrightarrow{\mu}\; L_{(y,y'')} \tag{2.7}$$

defined by $\mu((m,i,j;z),(m,j,k;z')) = (m,i,k;\, g_{ijk}(m)\, z\, z')$.

In general, $\mathcal{G} = (Y, B, L, \mu)$ is called an Abelian hermitian bundle gerbe with connection over M (in short: a **gerbe**), with curvature H, if $\pi : Y \to X$ is a surjective submersion, B is a 2-form on Y satisfying (2.5) called the gerbe's **curving** and L is a hermitian line bundle with connection over $Y^{[2]}$ satisfying (2.6) and equipped with a bilinear **groupoid multiplication** (2.7) preserving hermitian structure and parallel transport. In particular, the unity of the groupoid multiplication defines the canonical isomorphisms $L_{(y,y)} \cong \mathbb{C}$ and $L_{(y',y)} \cong L^*_{(y,y')}$. See [56] for a more formal definition. The most trivial example of a gerbe is the **unit gerbe** $(M, 0, M \times \mathbb{C}, \cdot\,)$ with vanishing curvature and with the groupoid structure in the trivial line bundle L over $M^{[2]} \cong M$ determined by the product in $\mathbb{C}$.

As we have seen, there is a gerbe corresponding to each family of local data. Conversely, given a gerbe $\mathcal{G}$, one may associate to it (non-canonically) a family of local data. In order to do this, one chooses local maps $\sigma_i : O_i \to Y$ such that $\pi \circ \sigma_i = id$ and local sections $s_{ij} : \sigma_{ij}(O_{ij}) \to L$, where $\sigma_{ij}(m) = (\sigma_i(m), \sigma_j(m))$, such that $s_{ji} \circ \sigma_{ji} = (s_{ij} \circ \sigma_{ij})^{-1}$. Then the relations

$$B_i = \sigma_i^* B\,, \tag{2.8,a}$$
$$\sigma_{ij}^*(\nabla s_{ij}) = \tfrac{1}{\mathrm{i}} A_{ij}\, s_{ij} \circ \sigma_{ij}\,, \tag{2.8,b}$$
$$\mu \circ (s_{ij} \circ \sigma_{ij} \otimes s_{jk} \circ \sigma_{jk}) = g_{ijk}\ s_{ik} \circ \sigma_{ik}\,. \tag{2.8,c}$$

determine gerbe local data (B_i, A_{ij}, g_{ijk}) with curvature H. Different choices of σ_i and s_{ij} lead to equivalent local data. We shall call the DD-class associated to such local data the DD-class of the gerbe.

Similarly, one may associate a geometric object to gerbe-module local data (Π_i, G_{ij}) corresponding to gerbe local data (B_i, A_{ij}, g_{ijk}). Again with $Y = \sqcup_i O_i$, one considers the N-dimensional hermitian vector bundle $E = \sqcup_i O_i \times \mathbb{C}^N$ over Y equipped with a (unitary) connection given over O_i by the local connection 1-forms Π_i. Let

$$\rho :\ L \otimes p_2^* E \ \longrightarrow\ p_1^* E$$

be the isomorphism of the vector bundles over $Y^{[2]}$ such that $\rho((m,i,j;z) \otimes (m,j;v)) = (m,i;G_{ij}(m)v)$. Isomorphism ρ is associative w.r.t. the groupoid multiplication μ:

$$\rho \circ (\mu \otimes id) \;=\; \rho \circ (id \otimes \rho)\,. \tag{2.9}$$

More generally, given a gerbe $\mathcal{G} = (Y, B, L, \mu)$, a pair $(E, \rho) = \mathcal{E}$ is called an N-dimensional hermitian $\mathcal{G}$-module with connection (in short: a **$\mathcal{G}$-module**) if E is an N-dimensional vector bundle with (unitary) connection over Y and ρ is a vector bundle isomorphism as above, preserving hermitian structure and connection and satisfying (2.9). Two $\mathcal{G}$-modules (E, ρ) and (E', ρ') are called **isomorphic** if there exists an isomorphism $\iota : E \to E'$ of hermitian vector bundles with connection such that

$$p_1^*\iota \circ \rho \;=\; \rho' \circ (id \otimes p_2^*\iota)\,.$$

The unit-gerbe modules coincide with hermitian vector bundles with connection over M.

As we have seen, there is a $\mathcal{G}$-module associated to a family (Π_i, G_{ij}) of gerbe-module local data corresponding to gerbe local data (B_i, A_{ij}, g_{ijk}), where gerbe $\mathcal{G}$ is constructed from the latter. Conversely, given a gerbe $\mathcal{G}$ and a $\mathcal{G}$-module $\mathcal{E} = (E, \rho)$, upon a choice of sections σ_i and s_{ij} for the former and of orthonormal bases (e_i^a) of sections of E over $\sigma_i(O_i)$, the relations

$$\sigma_i^* \nabla e_i^a \;=\; \tfrac{1}{\mathrm{i}}\, \Pi_i^{ba} \otimes e_i^b\,, \tag{2.10,a}$$
$$\rho \circ (s_{ij} \circ \sigma_{ij} \otimes e_j^a \circ \sigma_j) \;=\; G_{ij}^{ba}\, e_i^b \circ \sigma_i \tag{2.10,b}$$

define local gerbe-module data (Π_i, G_{ij}). Isomorphic $\mathcal{G}$-modules give rise to equivalent local data and conversely.

To reduce the structure group from $U(N)$ to a compact group G containing a central $U(1)$ subgroup, we may consider pairs $(P,\rho_{_P})$ where P is a principal G-bundle with connection over Y and

$$\rho_{_P}\,:\,SL\times_{Y^{[2]}}p_2^*P\ \longrightarrow\ p_1^*P$$

is an isomorphism of principal G-bundles with connection over $Y^{[2]}$, with SL denoting the circle subbundle of L. We assume that over $Y^{[3]}$,

$$\rho_{_P}\circ(\mu\times id)\ =\ \rho_{_P}\circ(id\times\rho_{_P})\,.$$

This implies that for each $y\in Y$, there is an action of $U(1)\cong SL_{(y,y)}$ on P_y and we shall require that it coincide with the restriction to $U(1)$ of the action of G on P_y. We shall call such pairs $(P,\rho_{_P})$ twisted G-bundles. Given an N-dimensional unitary representation U of G acting as identity on the $U(1)$ subgroup, the pair (E,ρ), where $E=P\times_G\mathbb{C}^N$ is the associated vector bundle and ρ is naturally induced by $\rho_{_P}$, forms a **$\mathcal{G}$-module associated to** $(P,\rho_{_P})$. We shall demand that the isomorphisms ι of the associated $\mathcal{G}$-modules be induced from the isomorphisms of the principal bundles that intertwine the maps $\rho_{_P}$. As for usual bundles, each N-dimensional gerbe module may be viewed as associated to its orthogonal frame bundle which is a twisted $U(N)$-bundle.

There are natural operations on gerbes and gerbe modules. One defines the **dual gerbe** $\mathcal{G}^*=(Y,-B,L^*,(\mu^*)^{-1})$ with curvature $-H$ and the **tensor product** of two gerbes

$$\mathcal{G}\otimes\mathcal{G}'=(Y\times_M Y',\,p^*B+p'^*B',\,(p^{[2]})^*L\otimes({p'}^{[2]})^*L',\,\mu\otimes\mu')$$

with curvature $H+H'$ where $p,\,p'$ stand for the projections of $Y\times_M Y'$ onto Y and Y', respectively. Note that the tensor product of $\mathcal{G}$ with the unit gerbe may be naturally identified with $\mathcal{G}$. For $f:N\to M$, $f^*\mathcal{G}$ will denote the **pullback gerbe** (with curvature f^*H) and for $N\subset M$, $\mathcal{G}|_N$ the **restriction** of $\mathcal{G}$ to N (the pullback by the embedding map). Similarly, for a $\mathcal{G}$-module $\mathcal{E}=(E,\rho)$, one may define a dual $\mathcal{G}^*$-module $\mathcal{E}^*=(E^*,(\rho^*)^{-1})$ as well as the pullbacks and the restrictions of $\mathcal{E}$. If $\mathcal{E}'=(E',\rho')$ is another $\mathcal{G}$-module then one may form the **direct sum** $\mathcal{G}$-module $\mathcal{E}\oplus\mathcal{E}'=(E\oplus E',\rho\oplus\rho')$. Moreover, if $\mathcal{E}'=(E',\rho')$ is a $\mathcal{G}'$-module, then $\mathcal{E}\otimes\mathcal{E}'=(p^*E\otimes p'^*E',\rho\otimes\rho')$ is a $(\mathcal{G}\otimes\mathcal{G}')$-module, the **tensor product** of $\mathcal{E}$ and $\mathcal{E}'$. If there exists a 1-dimensional $\mathcal{G}$-module $\mathcal{E}_1$ then any $\mathcal{G}$-module is isomorphic to a tensor product of $\mathcal{E}_1$ by a unit-gerbe module (i.e. by a vector bundle over M). We shall often use this property below.

Given two gerbes $\mathcal{G}$, $\mathcal{G}'$ over M, a 1-dimensional $(\mathcal{G}\otimes\mathcal{G}'^*)$-module $\mathcal{N}=(N,\nu)$ such that the curvature of the connection of N is equal to the 2-form $p'^*B'-p^*B$ on $Y\times_M Y'$ is called a **stable isomorphism** between $\mathcal{G}$ and $\mathcal{G}'$ and the gerbes for which such a module exists are said to be **stably isomorphic** [57]. Given a stable isomorphism between $\mathcal{G}$ and $\mathcal{G}'$, any $\mathcal{G}'$ module (E',ρ') induces in a natural way a $\mathcal{G}$-module (E,ρ) where $E_y=N_{(y,y')}\otimes E_{y'}$ for $(y,y')\in Y\times_M Y'$ (the latter spaces are identified for different y' with the use of the $\mathcal{G}\otimes\mathcal{G}'^*$ module structure on $\mathcal{N}$). One may this way compare gerbe modules over stably isomorphic gerbes. In particular, two gerbe modules induced by the same stable isomorphism are themselves isomorphic if and only if the original gerbe modules are.

Gerbes over M with curvature H exist if and only if the periods of H are in $2\pi\mathbb{Z}$. Two gerbes are stably isomorphic if and only if their local data are equivalent. For $H = 0$, the set of stable isomorphism classes forms a group under tensor product that may be identified with the cohomology group $H^2(M, U(1))$ by considering gerbes corresponding to local data $(0, 0, u_{ijk})$, where u_{ijk} is a $U(1)$-valued Čech 2-cocycle. Two stable isomorphism classes of gerbes with the same curvature differ by tensor multiplication by a class of zero curvature, i.e. by an element of $H^2(M, U(1))$.

$\mathcal{G}$-modules exist if and only if the DD-class of $\mathcal{G}$ is torsion i.e. if the curvature H of $\mathcal{G}$ is exact. Two $\mathcal{G}$-modules are isomorphic if and only if their local data associated to the same local data of $\mathcal{G}$ are equivalent.

3 Wess-Zumino action functional

Let Σ denote a 2-dimensional compact oriented surface and let M be a manifold. In 2d sigma models, Σ plays the role of a 2d spacetime or the string worldsheet and M that of the target space, with classical fields ϕ mapping Σ to M. In such models that appear in conformal field theory or in string theory, for example in the celebrated WZW model [71], one would like to consider the Wess-Zumino type contributions to the action functional of fields ϕ that may symbolically be written as

$$S_{WZ}(\phi) \;=\; \int\limits_{\Sigma} \phi^* d^{-1} H \tag{3.1}$$

where H is a closed 3-form on M. The problem with such topological[3] Wess-Zumino terms of the action is that the (Kalb-Ramond) 2-form $B = d^{-1}H$ such that $dB = H$ does not exist globally if H is not exact and even if it is, B is not unique. This is at this place that the gerbes and bundle gerbes show their utility [36, 46].

3.1 Closed string amplitudes

Let $\mathcal{G}$ be a gerbe with curvature H. Then $\phi^*\mathcal{G}$ is a gerbe over Σ (with curvature vanishing for dimensional reasons) and its stable equivalence class defines a cohomology class in $H^2(\Sigma, U(1))$. For Σ without boundary (such surfaces describe bulk 2d spacetime geometry or closed string worldsheets), $H^2(\Sigma, U(1)) = U(1)$ and one may define the closed string amplitude $\mathcal{A}(\phi) = \mathrm{e}^{\mathrm{i} S_{WZ}(\phi)}$ as equal to $\phi^*\mathcal{G}$ viewed as an element of $U(1)$ in a direct generalization of the notion of holonomy in line bundles [36], see also [60]. The amplitude defined this way depends only on the stable isomorphism class of $\mathcal{G}$. Given local data (B_i, A_{ij}, g_{ijk}) and a sufficiently fine triangulation of Σ with each triangle c contained in some subset O_{i_c}, one has [3, 36]

$$\mathcal{A}(\phi) \;=\; \exp\Big[i \sum_c \int\limits_c \phi^* B_{i_c} + i \sum_{b \subset c} \int\limits_b \phi^* A_{i_c i_b}\Big] \prod_{v \in b \subset c} g_{i_c i_b i_v}(\phi(v))\,, \tag{3.2}$$

where edges b and vertices v of the triangulation are contained in O_{i_b} and O_{i_v}, respectively. In the product over vertices on the right hand side, the convention is assumed that inverts the entry if the orientation of v inherited from c via b is negative

---

[3] i.e. independent of the metric on Σ

(i.e. if b is oriented from v to another vertex). This formula makes explicit the corrections needed to add to the sum of local contribution $\int\limits_c \phi^* B_{i_c}$ to the action for closed surfaces Σ.

3.2 Open string amplitudes: coupling to the Yang-Mills field

Surfaces with boundary describe 2d space-time geometry with boundaries or open-string worldsheets (with closed strings propagating in the handles). Suppose that $\partial\Sigma = \underset{s}{\sqcup}\, \ell^s$ where ℓ^s are the boundary loops that we shall, for convenience, parametrize by the standard circle S^1. Applying the same formula (3.2) to define the amplitude $\mathcal{A}(\phi)$ as in the closed string case, one observes that now the right hand side changes upon a change of the triangulation and the assignments i_b and i_v on the boundary. These changes may be summarized by saying that

$$\mathrm{e}^{\,\mathrm{i}S_{WZ}(\phi)} \;\in\; \underset{s}{\otimes}\, \mathcal{L}_{\phi|_{\ell^s}}$$

where $\mathcal{L}$ is a hermitian line bundle with connection over the loop space LM determined canonically by the local data (B_i, A_{ij}, g_{ijk}) of gerbe $\mathcal{G}$ [36]. In order to obtain number-valued open-string amplitudes, one may couple the ends of the open string to the (twisted) Yang-Mills field represented by a $\mathcal{G}$-module $\mathcal{E}$ with local data (Π_i, G_{ij}). The coupling is done by multiplying the amplitude $\mathrm{e}^{\,\mathrm{i}S_{WZ}(\phi)}$ by the traces of the appropriately defined holonomy of the boundary loops, see [46, 51, 17]. More precisely, define for a (pointed) loop $\varphi : S^1 \to M$ its holonomy by

$$\mathcal{H}(\varphi) \;=\; \mathcal{P} \prod_{v \in b} G_{i_v i_b}(\varphi(v)) \,\exp\Big[\, i \int\limits_b \varphi^* \Pi_{i_b} \Big] \tag{3.3}$$

for a sufficiently fine triangulation (split) of S^1 by intervals b with vertices v such that $\varphi(b) \subset O_{i_b}$ and $\varphi(v) \in O_{i_v}$. The operator $\mathcal{P}$ orders the terms and the exponetial path-wise (i.e. in agreement with the standard orientation of S^1) from right to left, starting from the vertex 1. Again, the matrix $G_{i_v i_b}$ should be inverted if v has negative orientation. Below, we shall use the same expression to define the parallel transport along open lines. For loops, let $\mathcal{W}(\phi) = \mathrm{tr}\, \mathcal{H}(\phi)$ be the corresponding Wilson loop "observable". Note that $\mathcal{W}(\varphi)$ does not change if we use in its definition local data (Π'_i, G'_{ij}) equivalent to (Π_i, G_{ij}). A straightforward check shows that under the change of the triangulation and its indexing, $\mathcal{W}(\varphi)$ transforms as an elements of $\mathcal{L}_\varphi^{-1}$. It follows that

$$\mathcal{A}(\phi) \;=\; \mathrm{e}^{\,\mathrm{i}S_{WZ}(\phi)} \prod_s \mathcal{W}(\phi|_{\ell^s})\,, \tag{3.4}$$

is independent of the choice of the indexed triangulation of Σ, provided that in the definition of $\mathcal{W}(\phi|_{\ell^s})$ one uses the indexed triangulations of the boundary loops induced from the indexed triangulation of Σ. The amplitude $\mathcal{A}(\phi)$ depends only on the stable isomorphism class of $\mathcal{G}$ and does not change if we use equivalent or induced local data for $\mathcal{E}$. If $\mathcal{E}$ is an associated gerbe module, then the holonomy $\mathcal{H}$ may be viewed as taking values in group G. We shall use then the notation $\mathcal{W}_U$ for the Wilson loop obtained by taking the trace of $\mathcal{H}$ in representation U of G.

3.3 Branes and general open string amplitudes

In the above definition of $\mathcal{A}(\phi)$ one may use holonomies with respect to different $\mathcal{G}$-modules for different boundary components. As we have seen above, existence of $\mathcal{G}$-modules puts a strong restriction on gerbe $\mathcal{G}$ requiring that its curvature be an exact form. Even if not satisfied globally, such a property might hold for the restriction of $\mathcal{G}$ to a submanifold of M. We shall call a pair $\mathcal{D} = (D, \mathcal{E})$, where D is a submanifold of M and $\mathcal{E}$ is an N-dimensional $\mathcal{G}|_D$-modules, a **$\mathcal{G}$-brane** (supported by D, of rank N). Two $\mathcal{G}$-branes will be called isomorphic if they have the same support D and if the corresponding $\mathcal{G}|_D$-modules are isomorphic. One may apply the above construction of amplitude $\mathcal{A}(\phi)$ if for each boundary component of Σ we are given a $\mathcal{G}$-brane $\mathcal{D}^s$ with support D^s, provided that we impose on maps ϕ the boundary conditions

$$\phi(\ell^s) \ \subset \ D^s \, . \tag{3.5}$$

We shall need to define amplitudes $\mathcal{A}(\phi)$ in a still more general situations [37, 39]. Let $\varphi \,:\, [0,\pi] \to M$ be an open line extending between submanifolds D^0 and D^1, i.e. such that

$$\varphi(0) \,\in\, D^0 \, , \qquad \varphi(\pi) \,\in\, D^1 \, .$$

We shall denote by $IM_{D^0}^{D^1}$ the space of such lines. Given N_s-dimensional $\mathcal{G}|_{D^s}$-modules $\mathcal{E}^s$, $s = 0, 1$, one may naturally define a hermitian vector bundle with connection $\mathcal{E}_{\mathcal{D}^0}^{\mathcal{D}^1}$ over $IM_{D^0}^{D^1}$ with typical fiber $\mathbb{C}^{N_0} \otimes \overline{\mathbb{C}^{N_1}}$. This is done in a way that is a straightforward generalization of the construction from [39] where the case of 1-dimensional modules has been treated, see Sects. 7 and 11.2 therein. Suppose now that the map $\phi : \Sigma \to M$ satisfies the boundary conditions (3.5) for ℓ^s being closed disjoint sub-intervals of the boundary loops of Σ, see Fig. 1.



Fig. 1

Then

$$\mathcal{A}(\phi) \ = \ \mathrm{e}^{\,\mathrm{i} S_{WZ}(\phi)} \, \Big(\prod_s{}' \, \mathcal{W}(\phi|_{\ell^s} \Big) \otimes \Big(\mathcal{P} \, \bigotimes_s{}'' \, \mathcal{H}(\phi|_{\ell^s}) \Big)$$

$$\in \quad \Big(\bigotimes_{m}\mathcal{L}_{\phi|_{\ell^m}}\Big)\otimes\Big(\mathcal{P}\bigotimes_{(s_-,s_+)}(\mathcal{E}^{\mathcal{D}^{s_+}}_{\mathcal{D}^{s_-}})_{\phi|_{\ell^{s_-s_+}}}\Big)\,, \tag{3.6}$$

where the product $\prod'$ is over ℓ^s that cover entirely a boundary component and the tensor product $\otimes''$ over the proper sub-intervals of boundary components, with the parallel transport $\mathcal{H}$ given by the same formula (3.3) as for closed loops. In the second line, ℓ^m run through the boundary loops without labeled sub-intervals and pairs (s_-,s_+) label subsequent pairs of proper sub-intervals of the boundary loops separated by unlabeled intervals $\ell^{s_-s_+}$. This is the higher rank version of the relation (7.11) of [39].

3.4 Purely classical approach

In the last incarnation, the topological amplitudes $\mathcal{A}(\phi)$, multiplied by the more standard non-topological[4] contributions, enter the functional integral expressions for general quantum amplitudes of the boundary two-dimensional field theory with the Wess-Zumino term (3.1) in the action. By themselves, they are, however, objects belonging to a mixed classical-quantum realm, the parallel transport or holonomy $\mathcal{H}$ being operator-valued, i.e. a quantum-mechanical concept. One may go back one step and obtain $\mathcal{A}(\phi)$ from entirely classical amplitudes. This discussion is somewhat parenthetical with respect to the main discourse of the paper and may be safely omitted by going directly to Sect. 4.

Let us first recall that the standard holonomy in the principal G-bundle P over M may be obtained by quantizing a mechanical system with the action $S_{_P}$ given by the connection form. More exactly, let φ be a loop in M and $t\mapsto\Phi(t)$ its lift to a loop in P. In a local trivialization where $P=O\times G$, $\Phi(t)=(\varphi(t),\gamma(t))$ and

$$S_{_P}(\Phi)\ =\ \int \mathrm{tr}\,\lambda\left(i\,\gamma^{-1}d\gamma\,+\,\gamma^{-1}(\varphi^*A)\,\gamma\right)\,,$$

where A is the local connection form with values in the Lie algebra $\mathbf{g}$ of G. Above, tr denotes a bilinear ad-invariant form on $\mathbf{g}$ and λ is a highest weight in the Cartan subalgebra $\mathbf{t}\subset\mathbf{g}$ defining a unitary representation U_λ of G. Note that the expression for the action is invariant under the local gauge transformations

$$\gamma\ \mapsto\ (\varphi^*h)^{-1}\gamma\,,\qquad A\ \mapsto\ h^{-1}A\,h\,+\,i\,h^{-1}dh$$

induced by a change of trivialization of P so that the action is globally defined. The Wilson loop $\mathcal{W}_{U_\lambda}(\varphi)$ in representation U_λ may be expressed as a path integral (that may be given a precise sense with the help of a Feynman-Kac type formula) over the lifts Φ:

$$\mathcal{W}_{U_\lambda}(\varphi)\ =\ \int \mathrm{e}^{\,\mathrm{i}S_P(\Phi)}\,D_\varphi\Phi$$

where, locally, the measure $D_\varphi\Phi=D\gamma$.

The same approach may be generalized to the case of Wilson loops in gerbe modules. Let $\mathcal{E}$ be a $\mathcal{G}$-module associated to a twisted G-bundle $(P,\rho_{_P})$. Observe that $P/U(1)$ descends to a G'-bundle P' over M for $G'=G/U(1)$. Let φ be a loop in M and Φ' its lift to a loop in P'. Choose local data (Π_i,G_{ij}) for $\mathcal{E}$. They provide local trivializations

---

[4] i.e. depending on the metric of Σ

$P'|_{O_i} \cong O_i \times G'$. For a triangulation of S^1 by intervals b with common vertices v such that $\varphi(b) \subset O_{i_b}$ and $\varphi(v) \in O_{i_v}$, choose maps $\gamma_b : b \mapsto G$ such that $(\varphi|_b, \gamma_b U(1))$ represent $\Phi'|_b$ in the local trivialization of $P'|_{O_{i_b}}$. Then for two subsequent intervals b_-, b_+ with common vertex v,

$$\gamma_{b_+}(v)\; u_v \;=\; G_{i_{b_+} i_v}(\varphi(v))\; G_{i_v i_{b_-}}(\varphi(v))\; \gamma_{b_-}(v)$$

for some $u_v \in U(1)$. Define

$$\mathrm{e}^{\,\mathsf{i}S_P(\Phi')} \;=\; \exp\Big[i\sum_b \int_b \mathrm{tr}\,\lambda\,\Big(i\,\gamma_b^{-1}d\gamma_b + \,\gamma_b^{-1}(\varphi^*\Pi_{i_b})\,\gamma_b\Big)\Big]\,\prod_v u_v\,,$$

for the highest weight λ satisfying $\mathrm{tr}\,\lambda\,\frac{1}{\mathsf{i}}\frac{d}{dt}\Big|_0 \mathrm{e}^{\mathsf{i}t} = 1$. It is easy to check that the above expression is independent of the choice of γ_b. On the other hand, under the change of indexed triangulations, it behaves as an element in $\mathcal{L}_\varphi^{-1}$. As for the untwisted case, the path integral of $\mathrm{e}^{\,\mathsf{i}S_P(\Phi')}$ over the lifts Φ' reproduces the Wilson loop $\mathcal{W}_{U_\lambda}(\varphi)$.

Let now $\phi : \Sigma \to M$ and (P^s, ρ_{P^s}) be twisted G^s-bundles. Then the purely classical amplitude

$$\mathcal{A}_{cl}\,(\phi, (\Phi'^s)) \;=\; \mathrm{e}^{\,\mathsf{i}S_{WZ}(\phi)}\,\prod_s \mathrm{e}^{\,\mathsf{i}S_{P^s}(\Phi'^s)}\,,$$

where Φ'^s are lifts to P'^s of the boundary loops $\phi|_{\ell^s}$, is unambiguously determined. If the twisted bundles (P^s, ρ_{P^s}) are defined only over submanifold $D^s \subset M$, one should impose the boundary conditions (3.5) on the map ϕ. The path integral over the lifts Φ'^s of the classical amplitudes $\mathcal{A}_{cl}\,(\phi, (\Phi'^s))$ reproduces the mixed amplitudes (3.4). It is not difficult to generate similar path-integral representations also for the general mixed amplitudes (3.6). Unlike the mixed formalism, the purely classical one lends itself to canonical quantization. This is the main reason why it is useful to consider it.

4 WZW models and gerbes over compact groups

In the WZW models of conformal field theory [71], the classical (Euclidean) fields ϕ map 2-dimensional Riemann surfaces into the manifold of a Lie group G that we shall assume here to be connected, compact and simple. The action functional is taken to be

$$S(\phi) \;=\; \tfrac{\mathsf{k}}{4\pi}\int_\Sigma \mathrm{tr}\,(\phi^{-1}\partial\phi)(\phi^{-1}\bar\partial\phi) \;+\; \int_\Sigma \phi^* d^{-1}H$$

where the closed 3-form H on G entering the Wess-Zumino term of the action is given by

$$H \;=\; \tfrac{\mathsf{k}}{12\pi}\,\mathrm{tr}\,(g^{-1}dg)^3\,. \tag{4.1}$$

The bilinear form tr on the Lie algebra $\mathbf{g}$ will be normalized so that for simply connected groups G the 3-form H has periods in $2\pi\mathbb{Z}$ if and only if k (called the level of the model) is an integer. As discussed in the previous section, in order to fully define the amplitudes $\mathrm{e}^{\,\mathsf{i}S(\phi)}$ for closed surfaces Σ we shall need a gerbe $\mathcal{G}_{\mathsf{k}}$ over G with curvature 3-form H.

4.1 Gerbes over simply connected groups

If G is a (connected, compact, simple) simply connected group, a gerbe over G with curvature H given by (4.1) exists if and only if k is an integer and it is unique up to stable isomorphisms since $H^2(G,U(1))=\{1\}$. Clearly one may take $\mathcal{G}_{\mathsf{k}}=\mathcal{G}_1^{\otimes\mathsf{k}}$. The basic gerbe $\mathcal{G}_1$ over simply connected groups G has been constructed in [53]. We shall describe gerbes $\mathcal{G}_{\mathsf{k}}$ (dropping below the subscript k) borrowing on the discussion in Sect. 2 of [40] and employing similar notations.

Let $\mathcal{A}_W$ denote the positive Weyl alcove in the Cartan subalgebra $\mathbf{t}$ of the Lie algebra $\mathbf{g}$ of G that will be identified with its dual $\mathbf{g}^*$ using the bilinear form defined by tr. The normalization of tr renders the length squared of the long roots equal to 2. $\mathcal{A}_W$ is a simplex with vertices τ_i, $i\in R\equiv\{0,1,\dots,r\}$, where r is the rank of $\mathbf{g}$. In particular, $\tau_0=0$ and $\tau_i=\frac{1}{k_i^\vee}\lambda_i$ for $i\neq 0$ with λ_i the simple weights and $k_i^\vee$ the dual Kac labels. For $i\in R$, let

$$\mathcal{A}_i=\{\,\tau\in\mathcal{A}_W\mid\tau=\sum_j s_j\tau_j \text{ with } s_i>0\,\},\qquad O_i=\{\,h\,\mathrm{e}^{2\pi\mathrm{i}\,\tau}\,h^{-1}\mid h\in G,\ \tau\in\mathcal{A}_i\,\}$$

and, for $I\subset R$, $\mathcal{A}_I=\underset{i\in I}{\cap}\mathcal{A}_i$ and $O_I=\underset{i\in I}{\cap}O_i$. Subsets $\mathcal{O}_i$ of G are open and $\mathcal{O}_I$ are composed of elements $h\,\mathrm{e}^{2\pi\mathrm{i}\,\tau}h^{-1}$ with $h\in G$ and $\tau\in\mathcal{A}_I$. The expressions

$$B_i\;=\;\tfrac{\mathsf{k}}{4\pi}\,\mathrm{tr}\,(h^{-1}dh)\,\mathrm{e}^{2\pi\mathrm{i}\tau}\,(h^{-1}dh)\,\mathrm{e}^{-2\pi\mathrm{i}\tau}\;+\;\mathrm{i}\mathsf{k}\,\mathrm{tr}\,(\tau-\tau_i)(h^{-1}dh)^2$$

define (smooth) 2-forms on O_i such that $dB_i=H|_{O_i}$. An important role is played by the subgroups

$$G_I\;=\;\{\,h\in G\mid h\,\mathrm{e}^{2\pi\mathrm{i}\,\tau}\,h^{-1}=\mathrm{e}^{2\pi\mathrm{i}\,\tau}\ \text{ for }\ \tau\in\mathcal{A}_I\setminus\underset{i\notin I}{\cup}\mathcal{A}_i\,\}$$

and the (smooth) maps

$$\mathcal{O}_I\ni g\;=\;h\,\mathrm{e}^{2\pi\mathrm{i}\tau}h^{-1}\;\xrightarrow{\rho_I}\;hG_I\in G/G_I$$

well defined because the adjoint action stabilizers of $\mathrm{e}^{2\pi\mathrm{i}\,\tau}$ for $\tau\in\mathcal{A}_I$ are contained in G_I. One introduces the principal G_I-bundles $\pi_I:P_I\to O_I$

$$P_I\;=\;\{\,(g,h)\in O_I\times G\mid\rho_I(g)=hG_I\,\}\,.$$

For the gerbes $\mathcal{G}=(Y,B,L,\mu)$, one sets

$$Y\;=\;\underset{i\in R}{\sqcup}P_i$$

with $\pi:Y\to G$ restricting to π_i on P_i and the 2-form B restricting to $\pi_i^*B_i$. Let

$$\widehat{Y}_{i_1..i_n}\;=\;P_I\times G_{i_1}\times\cdots\times G_{i_n}\qquad\text{and}\qquad Y_{i_1..i_n}\;=\;\widehat{Y}_{i_1..i_n}/G_I$$

for $I=\{i_1,\dots,i_n\}$ and G_I acting on $\widehat{Y}_{i_1..i_n}$ diagonally by the right multiplication. The fiber power $Y^{[n]}$ of Y may be identified with the disjoint union of $Y_{i_1..i_n}$ by assigning to the G_I-orbit of $((g,h),\gamma_1,..,\gamma_n)$ the n-tuple $(y_1,..,y_n)\in Y^{[n]}$ with $y_m=(g,h\gamma_m^{-1})$:

$$Y^{[n]}\;\cong\;\underset{(i_1,..,i_n)}{\sqcup}Y_{i_1..i_n}\,.$$

The construction of the line bundle L over $Y^{[2]}$ uses more detailed properties of the stabilizer groups G_I. For $I\subset J\subset R$, $G_I\supset G_J$ and the smallest of those groups, G_R, coincides with the Cartan subgroup T of G. Groups G_I are connected but not necessarily simply connected. Let $\mathbf{g}_I\supset\mathbf{t}$ denote the Lie algebra of G_I and let e_I be the exponential map from $\mathsf{i}\mathbf{g}_I$ to the universal cover $\widetilde{G}_I$. One has

$$G_I\ =\ \widetilde{G}_I/\mathcal{Z}_I\qquad\text{for}\qquad \mathcal{Z}_I\ =\ \mathrm{e}_I^{2\pi\mathsf{i}Q^\vee}$$

where $Q^\vee\subset\mathbf{t}$ is the coroot lattice of $\mathbf{g}$. Let $\chi_i:\mathcal{Z}_i\rightarrow U(1)$ be the character defined by

$$\chi_i(\mathrm{e}_i^{2\pi\mathsf{i}q})\ =\ \mathrm{e}^{2\pi\mathsf{i}\,\mathrm{tr}\,\tau_i q}\tag{4.2}$$

for $q\in Q^\vee$ and $\chi_{ij}:\widetilde{G}_{ij}\rightarrow U(1)$ be the 1-dimensional representation of $\widetilde{G}_{ij}$ given by the formula

$$\chi_{ij}(\widetilde{\gamma})\ =\ \exp\Big[\frac{1}{\mathsf{i}}\int\limits_{\widetilde{\gamma}}a_{ij}\Big]\tag{4.3}$$

where $a_{ij}\,=\,\mathsf{i}\,\mathrm{tr}\,(\tau_j-\tau_i)(\gamma^{-1}d\gamma)$ is a closed 1-form on G_{ij}. Let $\widehat{L}_{ij}$ denote the line bundle over $\widehat{Y}_{ij}$ with the fiber over $((g,h),\gamma,\gamma')$ composed of the equivalence classes $[\widetilde{\gamma},\widetilde{\gamma}',u]_{ij}$ with respect to the relation

$$(\widetilde{\gamma},\widetilde{\gamma}',u)\ \underset{ij}{\sim}\ (\widetilde{\gamma}\zeta,\widetilde{\gamma}'\zeta',\chi_i(\zeta)^{\mathsf{k}}\chi_j(\zeta')^{-\mathsf{k}}\,u)$$

for $\widetilde{\gamma}\in\widetilde{G}_i$, $\widetilde{\gamma}'\in\widetilde{G}_j$ projecting to $\gamma\in G_i$ and $\gamma'\in G_j$, respectively, and $u\in\mathbb{C}$, $\zeta\in\mathcal{Z}_i$, $\zeta'\in\mathcal{Z}_j$. We shall twist the flat structure of $\widehat{L}_{ij}$ by the connection form $A_{ij}=\mathsf{i}\mathsf{k}\,\mathrm{tr}\,(\tau_j-\tau_i)(h^{-1}dh)$. The right action of G_{ij} on $\widehat{Y}_{ij}$ lifts to the action on $\widehat{L}_{ij}$ defined by

$$((g,h),\,[\widetilde{\gamma},\widetilde{\gamma}',u]_{ij})\ \longmapsto\ ((g,h\gamma''),\,[\widetilde{\gamma}\widetilde{\gamma}'',\widetilde{\gamma}'\widetilde{\gamma}'',\chi_{ij}(\widetilde{\gamma}'')^{-\mathsf{k}}\,u]_{ij})$$

for $\gamma''\in G_{ij}$ and $\widetilde{\gamma}''$ its lift to $\widetilde{G}_{ij}$ (the right hand side is independent on the choice of the latter). The hermitian structure and the connection of $\widehat{L}_{ij}$ descend to the quotient bundle $\widehat{L}_{ij}/G_{ij}=L_{ij}$ over Y_{ij} and the line bundle L over $Y^{[2]}$ for the gerbe $\mathcal{G}$ is taken as equal to L_{ij} when restricted to Y_{ij}. Note that the curvature of L_{ij} is equal to the lift to Y_{ij} of the 2-form B_j-B_i on O_{ij}, as required. In general, unlike for the $SU(N)$ groups, there is no line bundle over O_{ij} with such curvature, hence the need of a more complicated construction of the gerbe [53].

The groupoid multiplication μ of $\mathcal{G}$ is defined as follows. Let $((g,h),\gamma,\gamma',\gamma'')\in\widehat{Y}_{ijk}$ represent $(y,y',y'')\in Y^{[3]}$ with $y=(g,h\gamma^{-1})$, $y'=(g,h\gamma'^{-1})$ and $y''=(g,h\gamma''^{-1})$ and let

$$\widehat{\ell}_{ij}\ =\ [\widetilde{\gamma},\widetilde{\gamma}',u]_{ij}\,,\qquad \widehat{\ell}_{jk}\ =\ [\widetilde{\gamma}',\widetilde{\gamma}'',u']_{jk}\,,\qquad \widehat{\ell}_{ik}\ =\ [\widetilde{\gamma},\widetilde{\gamma}'',uu']_{ij}$$

induce the elements $\ell_{ij}\in L_{(y,y')}$, $\ell_{jk}\in L_{(y',y'')}$ and $\ell_{ik}\in L_{(y,y'')}$. Then

$$\mu(\ell_{ij},\ell_{jk})\ =\ \ell_{ik}\,.$$

This ends the description of Meinrenken's gerbes $\mathcal{G}=(Y,B,L,\mu)$ over the simply connected group G.

4.2 Gerbes over non-simply connected groups

Let $G' = G/Z$, with Z a subgroup of the center of G, be a non-simply connected quotient of a simply connected group G. As mentioned in Introduction, to push down the gerbes $\mathcal{G}$ from G to G' one has to solve a cohomological equation. Let us start by identifying simple cohomological objects related to the pair (G, Z).

Choose for each $z \in Z$ an element $w_z \in G$ that normalizes the Cartan subgroup $T \subset G$ such that for $\tau \in \mathcal{A}_W$,

$$z\, e^{2\pi \mathsf{i} \tau} \;=\; w_z^{-1} \mathrm{e}^{2\pi \mathsf{i}\, z\tau}\, w_z \tag{4.4}$$

for some $z\tau \in \mathcal{A}_W$. For $z = 1$ we shall take $w_z = 1$. The map

$$\tau \;\longmapsto\; z\tau \;=\; w_z\, \tau\, w_z^{-1} \,+\, \tau_{z0}$$

is an affine transformation of $\mathcal{A}_W$ that permutes the vertices: $z\tau_i \equiv \tau_{zi}$. Let $b_{z,z'} \in \mathbf{t}$ be such that

$$w_z\, w_{z'}\, w_{zz'}^{-1} \;=\; c_{z,z'} \;=\; \mathrm{e}^{2\pi \mathsf{i}\, b_{z,z'}}\,. \tag{4.5}$$

We shall take $b_{z,1} = 0 = b_{1,z'}$. The Cartan subgroup valued chain $c = (c_{z,z'})$ is a 2-cocycle on group Z:

$$(\delta c)_{z,z'z''} \;=\; (w_z\, c_{z',z''} w_z^{-1})\, c_{zz',z''}^{-1}\, c_{z,z'z''}\, c_{z,z'}^{-1} \;=\; 1\,,$$

see Appendix A of [39] for a brief summary on finite group cohomology.

The 3-form H descends to a 3-form H' on G'. For (integer) k for which H' has periods in $2\pi\mathbb{Z}$ there exist a gerbe $\mathcal{G}' = (Y', B', L', \mu')$ over G' with curvature H'. To describe its explicit construction, we shall follow [40]. One takes $Y' = Y = \sqcup P_i$ with the natural projection π' on G' and $B' = B$. For $y = (g, h) \in P_i$ let zy denote the element $(zg, hw_z^{-1}) \in P_{zi}$. The fiber product space $Y'^{[n]}$ is composed of n-tuples $(y, y', \ldots, y^{(n-1)})$ such that for some $z, z', \ldots, z^{(n-2)} \in Z$

$$(y, zy', \cdot\cdot, z(z'(\cdot\cdot(z^{(n-2)}y^{(n-1)})\cdot\cdot))) \;\;\in\;\; Y^{[n]}\,.$$

One may then identify

$$Y'^{[n]} \;\cong \bigsqcup_{(z,z',\cdot\cdot,z^{(n-2)})\,\in\, Z^{n-1}} Y^{[n]} \;\cong \bigsqcup_{(z,z',\cdot\cdot,z^{(n-2)})\,\in\, Z^{n-1}}\; \bigsqcup_{(i_1,\cdot\cdot,i_n)} Y_{i_1..i_n}\,.$$

Let L' be the line bundle over $Y'^{[2]}$ that restricts to L on each component $Y^{[2]}$ in the above identification, i.e. to L_{ij} on $Y_{ij} \subset Y^{[2]}$. It remains to describe the groupoid multiplication μ'.

Let $(y, y', y'') \in Y'^{[3]}$ be such that $(y, zy', z(z'y'')) \in Y^{[3]}$. Identifying the triple $(y, zy', z(z'y''))$ with the G_{ijk}-orbit of $((g, h), \gamma, \gamma', \gamma'') \in \widehat{Y}_{ijk}$ for some (i, j, k) and employing the notations $i_z \equiv z^{-1}i$, $\gamma_z \equiv w_z^{-1}\gamma\, w_z \in G_{i_z}$ for $\gamma \in G_i$, we have

$$y \;=\; (g, h\gamma^{-1})\,, \qquad zy' \;=\; (g, h\gamma'^{-1})\,, \qquad z(z'y'') \;=\; (g, h\gamma''^{-1})\,, \tag{4.6}$$

$$y' \;=\; (z^{-1}g,\, h\, w_z \gamma_z'^{\,-1})\,, \qquad\qquad y'' \;=\; ((zz')^{-1}g,\, h\, w_z w_{z'} (\gamma''_z)^{-1}_{z'})\,, \tag{4.7}$$

$$z'y'' \;=\; (z^{-1}g,\, h\, w_z \gamma_z''^{\,-1})\,, \qquad (zz')y'' \;=\; (g,\, h\, (c^{-1}_{z,z'}\gamma'')^{-1})\,. \tag{4.8}$$

Recalling the explicit description of the line bundles L_{ij} as quotients of bundles $\widehat{L}_{ij}$, consider the equivalence classes

$$\widehat{\ell}_{ij} \;=\; [\widetilde{\gamma}, \widetilde{\gamma}', u]_{ij}\,, \quad \widehat{\ell}_{j_z k_z} \;=\; [\widetilde{\gamma}'_z, \widetilde{\gamma}''_z, u']_{j_z k_z}\,, \quad \widehat{\ell}_{ik} \;=\; [\widetilde{\gamma}, \widetilde{c}^{\,-1}_{z,z'} \widetilde{\gamma}'', u u']_{ik}\,, \tag{4.9}$$

with $\widetilde{\gamma} \in \widetilde{G}_i$ projecting to $\gamma \in G_i$ and $\widetilde{\gamma}_z \equiv w_z^{-1} \widetilde{\gamma}\, w_z \in \widetilde{G}_{i_z}$ etc. and with $\widetilde{c}_{z,z'} = \mathrm{e}_I^{2\pi \mathsf{i}\, b_{z,z'}} \in \widetilde{G}_I$. The above classes determine the elements

$$\ell_{ij} \in L_{(y,zy')} = L'_{(y,y')}\,, \quad \ell_{j_z k_z} \in L_{(y',z'y'')} = L'_{(y',y'')}\,, \quad \ell_{ik} \in L_{(y,(zz')y'')} = L'_{(y,y'')} \tag{4.10}$$

The groupoid multiplication μ' in L' is then defined by specifying the product of ℓ_{ij} and $\ell_{j_z k_z}$ via the relation [40]

$$\mu'(\ell_{ij},\, \ell_{j_z k_z}) \;=\; \mathrm{e}^{2\pi \mathsf{i} \mathsf{k}\, \mathrm{tr}\, \tau_k b_{z,z'}}\, V_{z,z'}\, \ell_{ik}\,, \tag{4.11}$$

where $V \;=\; (V_{z,z'})$ solves the cohomological equation

$$\delta V \;=\; U\,. \tag{4.12}$$

Explicitly, $(\delta V)_{z,z',z''} = V_{z',z''}\, V^{-1}_{zz',z''}\, V_{z,z'z''}\, V^{-1}_{z,z'}$ and $U = (U_{z,z',z''})$ is a 3-cocycle on Z with values in $U(1)$ defined by[5]

$$U_{z,z',z''} \;=\; \mathrm{e}^{-2\pi \mathsf{i} \mathsf{k}\, \mathrm{tr}\, \tau_{0_z}\, b_{z',z''}}$$

with $\tau_{0_z} = \tau_{z^{-1}0} = z^{-1}\tau_0$.

The cohomology class $[U] \in H^3(Z, U(1))$ of U does not depend on the ambiguity in the choice of w_z nor of $b_{z,z'}$. The changes

$$w_z \;\mapsto\; \mathrm{e}^{2\pi \mathsf{i}\, a_z} w_z\,, \qquad b_{z,z'} \;\mapsto\; b_{z,z'} \,+\, w_z a_{z'} w_z^{-1} \,-\, a_{zz'} \,+\, a_z \,+\, q_{z,z'} \tag{4.13}$$

with $a_z \in \mathbf{t}$ and $q_{z,z'} \in Q^\vee$ induces the transformation

$$U \;\longmapsto\; (\delta V')\, U \qquad \text{for} \qquad V'_{z,z'} \;=\; \mathrm{e}^{2\pi \mathsf{i} \mathsf{k}\, \mathrm{tr}\, \tau_{0_z}\, a_{z'}}\,.$$

The cohomological equation (4.12) has a solution only for k for which the cohomology class $[U]$ is trivial. The resulting set of levels k coincides with the one for which the periods of the 3-form H' are in $2\pi\mathbb{Z}$ [27, 40]. For all cases except when $G' = Spin(4n)/(\mathbb{Z}_2 \times \mathbb{Z}_2) = SO(4n)/\mathbb{Z}_2$ for $n = 2, 3, \ldots$, we have $H^2(Z, U(1)) = \{1\}$ and any two solutions V differ by a coboundary δW with $(\delta W)_{z,z'} = W_{z'} W^{-1}_{zz'} W_z$ and lead to stably isomorphic gerbes. In the exceptional case, $[U]$ is trivial if $\mathsf{k} \in \mathbb{Z}$ for n even and $\mathsf{k} \in 2\mathbb{Z}$ for n odd. In this case $H^2(\mathbb{Z}_2 \times \mathbb{Z}_2, U(1)) \cong \mathbb{Z}_2$ and there are two classes of non-equivalent solutions of Eq. (4.12) with all the solutions within one class differing by coboundaries δW. Such solutions lead to gerbes that fall into two different stable-isomorphism classes. In all cases one may choose $V_{z,z'}$ so that $V_{z,1} \equiv 1 \equiv V_{1,z'}$ which, in particular, assures that on $L'|_{Y^{[2]}} = L$ the groupoid product μ' coincides with μ.

---

[5] To obtain more concise expressions, we have multiplied the 3-cocycle U^{k} with U given by Eqs. (3.29) or (3.30) in [40] by the coboundary $\delta V'$ with $V'_{z,z'} = \mathrm{e}^{-2\pi \mathsf{i} \mathsf{k}\, \mathrm{tr}\, \tau_{zz'0}\, b_{z,z'}}$

5 Symmetric branes in the WZW models

In the open-string version of the WZW model, one may restrict the string ends to move on branes D in G and couple them to Yang-Mills fields on D. This is done using the concept of $\mathcal{G}$-branes as discussed in Sect. 3.2. One would like, however, to assure that the boundary version of the theory still possesses rich symmetry. The fundamental symmetry of the bulk WZW model is provided by two chiral copies of the current algebra $\widehat{\mathbf{g}}$ associated to the Lie algebra $\mathbf{g}$ of G. The simplest family of branes (called symmetric) assures that the diagonal current algebra is not broken in the presence of boundaries. Such requirement restricts the brane supports $D \subset G$ to coincide with a conjugacy class in G [2] and imposes further conditions on the $\mathcal{G}$-branes supported by D that we shall describe now. For simply connected G, the conjugacy classes may be labeled by the elements τ of the Weyl alcove $\mathcal{A}_W$ with

$$\mathcal{C}_\tau \;=\; \{\, h\,\mathrm{e}^{2\pi\mathrm{i}\tau}\, h^{-1} \mid h \in G \,\}\,.$$

being the class corresponding to τ. When restricted to $\mathcal{C}_\tau$, the 3-form H becomes exact. In particular, $H|_{\mathcal{C}_\tau} = dQ_\tau$ where

$$Q_\tau \;==\; \tfrac{\mathsf{k}}{4\pi}\,\mathrm{tr}\,(h^{-1}dh)\,\mathrm{e}^{2\pi\mathrm{i}\tau}(h^{-1}dh)\,\mathrm{e}^{-2\pi\mathrm{i}\tau}$$

is a (smooth) 2-form on $\mathcal{C}_\tau$. Let $\mathcal{G} = (Y, B, L, \mu)$ be a gerbe over G with curvature H as described above. Recall that a $\mathcal{G}$-brane $\mathcal{D}$ supported by D is a pair $(D, \mathcal{E})$ where $\mathcal{E}$ is a $\mathcal{G}|_D$-module. Such module determines, in turn, a vector bundle E with connection over $Y_D = \pi^{-1}(D)$ for π denoting the projection from Y to G. With $D = \mathcal{C}_\tau$, the additional restriction, imposed by the conservation of the diagonal current algebra, fixes the curvature of bundle E to be the scalar 2-form

$$F \;=\; \pi^*Q_\tau \;-\; B|_{Y_D}\,. \tag{5.1}$$

Note that this restricts very strongly the (twisted) Yang-Mills fields to which the ends of the string may be coupled without breaking the diagonal current-algebra symmetry.

The conjugacy classes in a non-simply connected group $G' = G/Z$ may in turn be labelled by the Z-orbits $[\tau]$ in the Weyl alcove $\mathcal{A}_W$ with

$$\mathcal{C}'_{[\tau]} \;=\; \mathcal{C}_\tau Z$$

for any $\tau \in [\tau]$. Let $Z_{[\tau]} \subset Z$ denotes the stabilizer subgroup of any $\tau \in [\tau]$. Note that $Z_{[\tau]}$ is composed of $z \in Z$ such that $z\mathcal{C}_\tau = \mathcal{C}_\tau$. It follows that

$$\mathcal{C}'_{[\tau]} \;\cong\; C_\tau / Z_{[\tau]}$$

for $\tau \in [\tau]$. In particular, $Z_{[\tau]}$ is the fundamental group of $\mathcal{C}'_{[\tau]}$ since the conjugacy classes $\mathcal{C}_\tau$ are simply connected. The 2-forms Q_τ project to a unique 2-form $Q'_{[\tau]}$ on $\mathcal{C}'_{[\tau]}$. If $\mathcal{G}' = (Y', B', L', \mu')$ is a gerbe over G' as described above then the symmetric $\mathcal{G}'$-branes supported by $D' = \mathcal{C}'_{[\tau]}$ are restricted by fixing the curvature of the vector bundle E' over $Y'_{D'} = \pi'^{-1}(\mathcal{C}'_{[\tau]})$ to be

$$F' \;=\; \pi'^*Q'_{[\tau]} \;-\; B'|_{Y'_{D'}}\,. \tag{5.2}$$

The main aim of this paper is to classify such symmetric $\mathcal{G}$- and $\mathcal{G}'$-branes up to isomorphism. For rank 1 $\mathcal{G}$-(or $\mathcal{G}'$-)branes, the curvature restriction may be reformulated by stating that the branes provide a stable isomorphism between the gerbes $\mathcal{G}|_D$ and $\mathcal{K} = (D, Q_\tau, D\times\mathbb{C}, \cdot\,)$, or between $\mathcal{G}'|_{D'}$ and $\mathcal{K}' = (D', Q'_{[\tau]}, D'\times\mathbb{C}, \cdot\,)$, and the notion of a rank 1 symmetric $\mathcal{G}$- or $\mathcal{G}'$-branes is equivalent to the one introduced in [39] and studied there in detail for groups $SU(N)$ and their non-simply connected quotients. Admitting rank N branes does not change much in the latter story, as we shall see below, but is necessary if we want to account properly for the conformal boundary conditions of general WZW models.

5.1 Symmetric branes for simply connected groups

Let $D = \mathcal{C}_\tau$ be the conjugacy class in the simply connected group G and Y_D be defined as above. Clearly

$$Y_D \;=\; \sqcup_i P_i|_{\mathcal{C}_\tau} \tag{5.3}$$

where $P_i|_{\mathcal{C}_\tau}$ is non-empty if and only if $\tau\in\mathcal{A}_i$. Let G_τ denote the adjoint-action isotropy subgroup of $\mathrm{e}^{2\pi\mathrm{i}\tau}$ so that $\mathcal{C}_\tau \cong G/G_\tau$. G_τ is connected and $T\subset G_\tau\subset G_i$ if $\tau\in\mathcal{A}_i$ (G_τ is is one of groups G_I with $I\ni i$). We shall denote by $\mathbf{g}_\tau$ the Lie algebra of G_τ. The space

$$P_\tau \;=\; \{\,(g,h)\in\mathcal{C}_\tau\times G \mid g = h\,\mathrm{e}^{2\pi\mathrm{i}\tau}\,h^{-1}\,\}$$

equipped with the projection on the first factor and the right action of G_τ on the second factor becomes a principal G_τ-bundle over $\mathcal{C}_\tau$. Of course, $P_\tau\cong G$. Note that if $\tau\in\mathcal{A}_i$ then $P_\tau\subset P_i|_{\mathcal{C}_\tau}$. Besides, we may identify

$$P_i|_{\mathcal{C}_\tau} \;\cong\; Y_{i\tau} \;=\; \widehat{Y}_{i\tau}/G_\tau \qquad\text{for}\qquad \widehat{Y}_{i\tau} \;=\; P_\tau\times G_i \tag{5.4}$$

assigning to the orbit of $((g,h),\gamma)\in\widehat{Y}_{i\tau}$ under the right diagonal action of G_τ the element $(g,h\gamma^{-1})\in P_i$.

Let us start by describing the rank 1 symmetric $\mathcal{G}$-branes $(\mathcal{C}_\tau,\mathcal{E}_1)$ supported by the conjugacy classes $\mathcal{C}_\tau$ such that $\mathsf{k}\tau=\lambda$ is a weight. Weights $\lambda=\mathsf{k}\tau$ for $\tau\in\mathcal{A}_W$ are often called integrable [45] (at level k) and we shall also call the corresponding conjugacy classes **integrable**. The construction of the line bundle E over Y_D such that $\mathcal{E}_1=(E,\rho)$ will resemble that of the line bundle L in Sect. 4.1. Let $\widetilde{G}_\tau$ denote the universal covering group of $G_\tau=\widetilde{G}_\tau/\mathcal{Z}_\tau$ where $\mathcal{Z}_\tau=\mathrm{e}_\tau^{2\pi\mathrm{i}Q^\vee}$ for e_τ standing for the exponential map from $\mathrm{i}\mathbf{g}_\tau$ to $\widetilde{G}_\tau$. Consider the 1-form $a_{i\tau}=\mathrm{i}\,\mathrm{tr}\,(\tau-\tau_i)(\gamma^{-1}d\gamma)$ on G_τ. It is easy to see that the adjoint action of G_τ preserves $\tau-\tau_i$. As a result, $a_{i\tau}$ is a closed form. By the formula

$$\chi_{i\tau}(\widetilde{\gamma}) \;=\; \exp\Big[\tfrac{1}{\mathrm{i}}\int\limits_{\widetilde{\gamma}} a_{i\tau}\Big]$$

it defines a 1-dimensional representation $\chi_{i\tau}:\widetilde{G}_\tau\to U(1)$, compare (4.3). The embedding $\mathbf{g}_\tau\subset\mathbf{g}_i$ induces the canonical homomorphism $\widetilde{G}_\tau\to\widetilde{G}_i$ that maps $\mathcal{Z}_\tau$ to $\mathcal{Z}_i$ sending $\zeta_\tau=\mathrm{e}_\tau^{2\pi\mathrm{i}q}\in\mathcal{Z}_\tau$ to $\zeta_i=\mathrm{e}_i^{2\pi\mathrm{i}q}\in\mathcal{Z}_i$ for $q\in Q^\vee$. Note that

$$\chi_{i\tau}(\widetilde{\gamma}\zeta_\tau)^{\mathsf{k}} \;=\; \chi_{i\tau}(\widetilde{\gamma})^{\mathsf{k}}\,\chi_i(\zeta_i)^{-\mathsf{k}}\,\mathrm{e}^{2\pi\mathrm{i}\,\mathsf{k}\,\mathrm{tr}\,\tau q}$$

and the last factor drops out if $\mathsf{k}\tau$ is a weight.

Let $\widehat{L}_i$ denote the flat line bundle over G_i with the fiber over γ composed of the equivalence classes $[\widetilde{\gamma},u]_i$ with respect to the equivalence relation

$$(\widetilde{\gamma},v) \ \underset{i}{\sim}\ (\widetilde{\gamma}\zeta,\chi_i(\zeta)^{\mathsf{k}}v) \tag{5.5}$$

for $\widetilde{\gamma}\in\widetilde{G}_i$ projecting to γ, $\zeta\in\mathcal{Z}_i$ and $v\in\mathbb{C}$, see (4.2). We shall denote by $\widehat{L}_{i\tau}$ the pullback of $\widehat{L}_i$ to $\widehat{Y}_{i\tau}=P_\tau\times G_i$ with the flat structure twisted by the connection form $A_{i\tau}\,=\,\mathsf{i}\mathsf{k}\,\mathrm{tr}\,(\tau-\tau_i)(h^{-1}dh)$. The right action of G_τ on $\widehat{Y}_{i\tau}$ lifts to the action on $\widehat{L}_{i\tau}$ defined by

$$((g,h),\,[\widetilde{\gamma},v]_i) \ \longmapsto\ ((g,h\gamma_\tau),\,[\widetilde{\gamma}\widetilde{\gamma}_\tau,\chi_{i\tau}(\widetilde{\gamma}_\tau)^{-\mathsf{k}}v]_i) \tag{5.6}$$

for $\widetilde{\gamma}_\tau\in\widetilde{G}_\tau$ projecting to $\gamma_\tau\in G_\tau$. For $\mathsf{k}\tau$ a weight, the right hand side does not depend on the choice of the lift $\widetilde{\gamma}_\tau$. The hermitian structure and the connection on $\widehat{L}_{i\tau}$ descend to the quotient bundle $\widehat{L}_{i\tau}/G_\tau=L_{i\tau}$ over $Y_{i\tau}$. The curvature form $F_{i\tau}$ of $L_{i\tau}$ is given by the relation:

$$\widehat{\iota}^*F_{i\tau} \ =\ i\mathsf{k}\,\mathrm{tr}\,(\tau_i-\tau)(h^{-1}dh)^2$$

with $\widehat{\iota}$ standing for the projection from $\widehat{Y}_{i\tau}$ to $Y_{i\tau}$. Let E be a hermitian line bundle with connection over Y_D coinciding with $L_{i\tau}$ over $Y_{i\tau}$, see (5.3) and (5.4). Note that E satisfies the curvature constraint (5.1), as required. Besides, there exist a bundle isomorphism

$$\rho:\ L|_{Y_D^{[2]}}\otimes p_2^*E \ \longrightarrow\ p_1^*E \tag{5.7}$$

satisfying relation (2.9). It is defined in the following way. Let, for $\tau\in\mathcal{A}_{ij}$, $(g,h)\in P_\tau$, $\widetilde{\gamma}\in\widetilde{G}_i$ projecting to $\gamma\in G_i$ and $\widetilde{\gamma}'\in\widetilde{G}_j$ projecting to $\gamma'\in G_j$,

$$\widehat{\ell}_{ij} \ =\ [\widetilde{\gamma},\widetilde{\gamma}',u]_{ij}\,,\qquad \widehat{\ell}_{i\tau}\ =\ [\widetilde{\gamma},uv']_i\,,\qquad \widehat{\ell}_{j\tau}\ =[\widetilde{\gamma}',v']_{j\tau}$$

induce, respectively, the elements $\ell_{ij}\in L_{(y,y')}$, $\ell_{i\tau}\in E_y$ and $\ell_{j\tau}\in E_{y'}$ for $y=(g,h\gamma^{-1})$ and $y'=(g,h\gamma'^{-1})$. Then

$$\rho(\ell_{ij}\otimes\ell_{j\tau}) \ =\ \ell_{i\tau}\,.$$

We infer that if $\mathsf{k}\tau$ is a weight then (E,ρ) defines a 1-dimensional $\mathcal{G}|_D$-module $\mathcal{E}_1$. Besides, the curvature of bundle E satisfies the constraint (5.1). Consequently, $\mathcal{D}_1=(\mathcal{C}_\tau,\mathcal{E}_1)$ is a symmetric rank 1 $\mathcal{G}$-brane supported by $D=\mathcal{C}_\tau$. General symmetric $\mathcal{G}$-branes supported by the integrable conjugacy classes are isomorphic to the ones obtained by tensoring E with a trivial N-dimensional bundle (all flat vector bundles over $\mathcal{C}_\tau$ are trivial up to isomorphism). In other words, they correspond to rank N $\mathcal{G}|_D$-modules

$$\mathcal{E}_N \ =\ \mathcal{E}_1\underset{N\ \text{terms}}{\oplus\ldots\oplus}\mathcal{E}_1\ .$$

A rank N $\mathcal{G}$-brane $\mathcal{D}_N=(\mathcal{C}_\tau,\mathcal{E}_N)$ obtained this way is called a stack of N $\mathcal{G}$-branes $\mathcal{D}_1$ in physicists' language.

The non-integrable conjugacy classes $D=\mathcal{C}_\tau$, i.e. such that $\mathsf{k}\tau$ is not a weight, support no symmetric $\mathcal{G}$-branes, as we shall show now. Existence of such a brane implies that $Y|_D$ admits a vector bundle E with curvature F given by (5.1), i.e. restricting on each $Y_{i\tau}$ to the scalar 2-form $F_{i\tau}$. The periods of $F_{i\tau}$ over 2-spheres in $Y_{i\tau}$ have then to belong to $2\pi\mathbb{Z}$. Let $\alpha_j^\vee$ for $j=1,\dots,r$ be the $j^{\rm th}$ simple coroot of $\mathbf{g}$ and $e_{\pm\alpha_j}$ be the corresponding step generators. Consider for each $j\neq i$ the $su(2)$ subalgebra of $\mathbf{g}$ which is the real form of the subalgebra of $\mathbf{g}^{\mathbb{C}}$ generated by $\alpha_j^\vee$ and $e_{\pm\alpha_j}$. Such an $su(2)$ subalgebra exponentiates to a subgroup $SU(2)\subset G_i\subset G$ whose diagonal embedding into $G\times G_i\cong\widehat{Y}_{i\tau}$ induces the map from $S^2\cong SU(2)/U(1)$ to $Y_{i\tau}$. The period of $F_{i\tau}$ on such a 2-cycle is equal to

$$2\pi\mathsf{k}\,\mathrm{tr}(\tau-\tau_i)\alpha_j^\vee\;=\;2\pi\mathsf{k}\,\mathrm{tr}\,\tau\alpha_j^\vee\,.$$

If τ is not equal to a vertex of the Weyl alcove $\mathcal{A}_W$ then $Y_{i\tau}$ is not empty (i.e. $\tau\in\mathcal{A}_i$) for at least two values of i and we infer that $\mathsf{k}\tau$ must be a weight. If τ is a vertex of $\mathcal{A}_W$ then it belongs to the unique $\mathcal{A}_i$ for which $\tau=\tau_i$ and the 2-form $F_{i\tau}$ vanishes. In this case $Y_D\cong P_{\tau_i}\cong G$ and $\mathcal{G}$-brane's vector bundle E over Y_D must be flat and hence isomorphic to a trivial N-dimensional bundle since G is simply connected. On the other hand, $Y_D^{[2]}\cong\{\,(h\gamma^{-1},h)\mid h\in G\,,\ \gamma\in G_i\,\}\cong G\times G_i$ with the line bundle $L|_{Y_D^{[2]}}$ isomorphic to the pullback of $\widehat{L}_i$ from G_i. The isomorphism (5.7) maps now flat bundles and is given by a constant map $\mathsf{R}:G\times\widetilde{G}_i\longrightarrow GL(N)$ such that

$$\mathsf{R}(h,\widetilde{\gamma}\zeta)\;=\;\chi_i(\zeta)^{-\mathsf{k}}\,\mathsf{R}(h,\widetilde{\gamma})\,.$$

Clearly, such a constant map exists only if $\mathsf{k}\tau_i$ is a weight so that the right hand side is ζ-independent.

In summary, in the case of simply connected groups, the symmetric $\mathcal{G}$-branes are determined up to isomorphism by giving their support, an integrable conjugacy class $\mathcal{C}_\tau$, and their rank N.

5.2 Symmetric branes for non-simply connected groups

Let $G'=G/Z$ be a non-simply connected group and $\mathcal{G}'=(Y',B',L',\mu')$ be a gerbe over it described in Sect. 4.2. We shall look here for symmetric $\mathcal{G}'$-branes supported by the conjugacy classes $\mathcal{C}'_{[\tau]}\subset G'$. This again leads to a problem in finite group Z cohomology, as already noted in [39] for the quotient groups of $SU(N)$. For $D'=\mathcal{C}'_{[\tau]}$,

$$Y'_{D'}\;=\;\sqcup_{\tau\in[\tau]}\sqcup_{i}P_i|_{\mathcal{C}_\tau}\,.\tag{5.8}$$

Any symmetric $\mathcal{G}'$-brane $\mathcal{D}'=(D',\mathcal{E}')$ supported by $\mathcal{C}'_{[\tau]}$ induces by restriction symmetric $\mathcal{G}$-branes supported by $\mathcal{C}_\tau$ for $\tau\in[\tau]$ (note that $Y_D\subset Y'_{D'}$ for $D=\mathcal{C}_\tau$ etc.). It follows that only integrable conjugacy classes $\mathcal{C}'_{[\tau]}$ for which $\mathsf{k}\tau$ is a weight may support symmetric $\mathcal{G}'$-branes (if $\mathsf{k}\tau$ is a weight for some $\tau\in[\tau]$ then it is for all $\tau\in[\tau]$). Since now on we shall then assume that $\mathsf{k}\tau$ is a weight for $\tau\in[\tau]$.

Let us first look for 1-dimensional $\mathcal{G}'_{D'}$-modules $\mathcal{E}'_1=(E',\rho')$ satisfying the curvature constraint (5.2). By the above argument, we may take E' as the line bundle over $Y'_{D'}$

that over each $P_i|_{\mathcal{C}_\tau} \cong Y_{i\tau}$ restricts to $L_{i\tau}$. The problem is to define the isomorphism

$$\rho' : \, L'|_{Y'^{[2]}_{D'}} \otimes p_2^* E' \;\longrightarrow\; p_1^* E' \tag{5.9}$$

that satisfies the associativity condition

$$\rho' \circ (\mu' \otimes id) \;=\; \rho' \circ (id \otimes \rho')\,, \tag{5.10}$$

see (2.9). Let $(y, y') \in Y'^{[2]}_{D'}$. This means that, in the notations of Sect. 4.2,

$$y = (g, h\gamma^{-1})\,, \qquad zy' = (g, h\gamma'^{-1})\,, \qquad y' = (z^{-1}g, hw_z\gamma_z'^{-1})$$

with $g = h\mathrm{e}^{2\pi\mathrm{i}\tau}h^{-1}$ for some $\tau \in [\tau] \cap \mathcal{A}_{ij}$, $\gamma \in G_i$, $\gamma' \in G_j$ and $z \in Z$. The equivalence classes

$$\widehat{\ell}_{ij} \;=\; [\widetilde{\gamma}, \widetilde{\gamma}', u]_{ij}\,, \qquad \widehat{\ell}_{i\tau} \;=\; [\widetilde{\gamma}, uv']_i\,, \qquad \widehat{\ell}_{j_z\tau_z} \;= [\widetilde{\gamma}'_z, v']_{j_z}\,, \tag{5.11}$$

with $\tau_z \equiv z^{-1}\tau$ and with $\widetilde{\gamma} \in \widetilde{G}_i$ projecting to γ and $\widetilde{\gamma}' \in \widetilde{G}_j$ to γ', induce, respectively, the elements $\ell_{ij} \in L_{(y,zy')} = L'_{(y,y')}$, $\ell_{i\tau} \in E'_y$ and $\ell_{j_z\tau_z} \in E'_{y'}$. Since the isomorphism ρ' is required to preserve the connection, necessarily,

$$\rho'(\ell_{ij} \otimes \ell_{j_z\tau_z}) \;=\; \mathcal{W}^{ij}_{\tau;z}\,\ell_{i\tau} \tag{5.12}$$

for $\mathcal{W}^{ij}_{\tau;z} \in U(1)$. Besides, the conjugation of ρ' with an isomorphism of E' reducing to multiplication by $\mathcal{Y}^i_\tau \in U(1)$ over the connected components $P_i|_{\mathcal{C}_\tau}$ of $Y'_{D'}$, results in the change

$$\mathcal{W}^{ij}_{\tau;z} \;\longmapsto\; \mathcal{Y}^i_\tau\,\mathcal{W}^{ij}_{\tau;z}\,\mathcal{Y}^{j_z\,-1}_{\tau_z}\,. \tag{5.13}$$

Such changes lead to isomorphic $\mathcal{G}'|_{D'}$-modules and hence to isomorphic $\mathcal{G}'$-branes.

It is now easy to translate identity (5.10) to conditions for coefficients $\mathcal{W}^{ij}_{\tau;z}$. For a triple $(y, y', y'') \in Y'^{[3]}_{D'}$ as in (4.6-4.8) where $g = h\mathrm{e}^{2\pi\mathrm{i}\tau}h^{-1}$ for some $\tau \in [\tau] \cap \mathcal{A}_{ijk}$, $\gamma \in G_i$, $\gamma' \in G_j$, $\gamma'' \in G_k$ and $z, z' \in Z$, consider the equivalence classes $\widehat{\ell}_{ij}$, $\widehat{\ell}_{j_zk_z}$ and $\widehat{\ell}_{ik}$ generating elements ℓ_{ij}, $\ell_{j_zk_z}$ and ℓ_{ik} as in (4.9) and (4.10). Besides, let

$$\widehat{\ell}_{k_{zz'}\tau_{zz'}} = [(\widetilde{c}^{\,-1}_{z,z'}\widetilde{\gamma}'')_{zz'}, v'']_{k_{zz'}}\,, \qquad \widehat{\ell}'_{k_{zz'}\tau_{zz'}} = [(\widetilde{\gamma}''_z)_{z'}, \chi_{k\tau}(\widetilde{c}_{z,z'})^{-\mathsf{k}}v'']_{k_{zz'}}\,. \tag{5.14}$$

It is easy to check employing the action (5.6) of $G_{\tau_{zz'}}$ and the relations

$$(\widetilde{\gamma}''_z)_{z'} = (\widetilde{c}^{\,-1}_{z,z'}\widetilde{\gamma}''\widetilde{c}_{z,z'})_{zz'}\,, \qquad \chi_{k_{zz'}\tau_{zz'}}((\widetilde{c}_{z,z'})_{zz'}) = \chi_{k\tau}(\widetilde{c}_{z,z'})$$

that both equivalence classes define the same element $\ell_{k_{zz'}\tau_{zz'}} \in E'_{y''}$. Using Eqs. (4.11) and (5.12) and the first way of (5.14) to represent $\ell_{k_{zz'}\tau_{zz'}}$, we infer that

$$\begin{aligned}\rho'(\mu'(\ell_{ij}, \ell_{j_zk_z}) \otimes \ell_{k_{zz'}\tau_{zz'}}) \;&=\; \mathrm{e}^{2\pi\mathrm{i}\mathsf{k}\,\mathrm{tr}\,\tau_k b_{z,z'}}\,V_{z,z'}\;\rho'(\ell_{ik} \otimes \ell_{k_{zz'}\tau_{zz'}})\\ &=\; \mathrm{e}^{2\pi\mathrm{i}\mathsf{k}\,\mathrm{tr}\,\tau_k b_{z,z'}}\,V_{z,z'}\,\mathcal{W}^{ik}_{\tau;zz'}\;\ell_{i\tau}\end{aligned}$$

if $v' = u'v''$ in (5.11). On the other hand, using the second way to represent $\ell_{k_{zz'}\tau_{zz'}}$, we obtain

$$\rho'(\ell_{ij} \otimes \rho'(\ell_{j_zk_z} \otimes \ell_{k_{zz'}\tau_{zz'}})) \;=\; \chi_{k\tau}(\widetilde{c}_{z,z'})^{-\mathsf{k}}\;\rho'(\ell_{ij} \otimes \mathcal{W}^{j_zk_z}_{\tau_z;z'}\,\ell_{j_z\tau_z})$$

$$= \ \chi_{k\tau}(\widetilde{c}_{z,z'})^{-\mathsf{k}}\, \mathcal{W}^{ij}_{\tau;z}\, \mathcal{W}^{j_zk_z}_{\tau_z;z'}\ \ell_{i\tau}\,.$$

The associativity condition (5.10) requires that both sides be equal and, since $\chi_{k\tau}(\widetilde{c}_{z,z'}) = \mathrm{e}^{2\pi\mathrm{i}\mathsf{k}\,\mathrm{tr}\,(\tau-\tau_k)b_{z,z'}}$, it is equivalent to the identity

$$\mathcal{W}^{j_zk_z}_{\tau_z;z'}\, \mathcal{W}^{ik\ -1}_{\tau;zz'}\, \mathcal{W}^{ij}_{\tau;z} \ = \ \mathcal{V}_{\tau;z,z'} \tag{5.15}$$

with

$$\mathcal{V}_{\tau;z,z'} \ = \ \mathrm{e}^{2\pi\mathrm{i}\mathsf{k}\,\mathrm{tr}\,\tau b_{z,z'}}\, V_{z,z'} \tag{5.16}$$

where $V_{z,z'}$ is the solution of Eq. (4.12) that determines gerbe $\mathcal{G}'$. It is easy to verify that $\mathcal{V} = (\mathcal{V}_{\tau;z,z'})$ forms a 2-cocycle on group Z with values in the Z-module $U(1)^{[\tau]}$ of $U(1)$-valued functions on the Z-orbit $[\tau]$, i.e. that

$$(\delta\mathcal{V})_{\tau;z,z',z''} \ = \ \mathcal{V}_{\tau_z;z',z''}\, \mathcal{V}^{-1}_{\tau;zz',z''}\, \mathcal{V}_{\tau;z,z'z''}\, \mathcal{V}^{-1}_{\tau;z,z'} \ = \ 1$$

for $\tau \in [\tau]$ and $z, z', z'' \in Z$. The cohomology class $[\mathcal{V}] \in H^2(Z, U(1)^{[\tau]})$ is independent of the choices of $b_{z,z'}$. Indeed, under the transformations (4.13),

$$\mathcal{V} \ \longmapsto (\delta\mathcal{W}')\, \mathcal{V} \qquad \text{for} \qquad \mathcal{W}'_{\tau;z} \ = \ \mathrm{e}^{2\pi\mathrm{i}\mathsf{k}\,\mathrm{tr}\,\tau a_z}\,. \tag{5.17}$$

Let us first show that if Eq. (5.15) has a solution then by a transformation (5.13), which maps solutions to solutions, one may achieve that $\mathcal{W}^{ij}_{\tau;z}$ does not depend on (i,j). We shall use the fact that $\mathcal{V}_{\tau;z,1} = \mathcal{V}_{\tau;1,z'} \equiv 1$. For $z, z' = 1$, Eq. (5.15) reduces to

$$\mathcal{W}^{jk}_{\tau;1}\, \mathcal{W}^{ik\,-1}_{\tau;1}\, \mathcal{W}^{ij}_{\tau;1} \ = \ 1$$

which means that, for each τ, $(\mathcal{W}^{ij}_{\tau;1})$ is a $U(1)$-valued Čech 1-cocycle for the covering $(\mathcal{A}_i)$ of the point $\{\tau\}$. Such a 1-cocycle is necessarily a coboundary:

$$\mathcal{W}^{ij}_{\tau;1} \ = \ \mathcal{Y}^{i\ -1}_{\tau}\, \mathcal{Y}^{j}_{\tau}$$

for $\mathcal{Y}^i_\tau \in U(1)$. By the transformation (5.13), we assure that $\mathcal{W}^{ij}_{\tau;1} = 1$. Now, specifying Eq. (5.15) to $z = 1$ or to $z' = 1$, we infer that

$$\mathcal{W}^{jk}_{\tau;z'} \ = \ \mathcal{W}^{ik}_{\tau;z'}\,, \qquad \mathcal{W}^{ij}_{\tau;z} \ = \ \mathcal{W}^{ik}_{\tau;z}\,,$$

i.e. that $\mathcal{W}^{ij}_{\tau;z}$ is independent of (i,j). Note in passing that the condition $\mathcal{W}^{ij}_{\tau;1} = 1$ means that the isomorphism ρ' restricts to ρ over $Y^{[2]}_D \subset Y'^{[2]}_{D'}$ for $D = \mathcal{C}_\tau$ with $\tau \subset [\tau]$.

Without the (i,j)-dependence, $(\mathcal{W}_{\tau;z}) \equiv \mathcal{W}$ forms a 1-cochain on group Z with values in the Z-module $U(1)^{[\tau]}$. The identity (5.15) reduces then to the cohomological equation

$$\delta\mathcal{W} \ = \ \mathcal{V} \tag{5.18}$$

where $(\delta\mathcal{W})_{\tau;z,z'} \ = \ \mathcal{W}_{\tau_z;z'}\, \mathcal{W}^{-1}_{\tau;zz'}\, \mathcal{W}_{\tau;z}$. Eq. (5.18) has a solution if and only if the cohomology class

$$[\mathcal{V}] \ \in \ H^2(Z, U(1)^{[\tau]}) \ \cong \ H^2(Z_{[\tau]}, U(1))$$

is trivial where, as above, $Z_{[\tau]} \subset Z$ denotes the stabilizer subgroup of $\tau \in [\tau]$. Vanishing of the cohomology class $[\mathcal{V}]$ is a necessary and sufficient condition for the existence of a rank 1 symmetric $\mathcal{G}'$-brane supported by the integrable conjugacy class $\mathcal{C}'_{[\tau]}$. Two solutions of Eq. (5.18) lead to isomorphic $\mathcal{G}'$-branes if and only if they differ by a coboundary $\delta\mathcal{Y}$ with $(\delta\mathcal{Y})_{\tau;z} = \mathcal{Y}_{\tau_z}\mathcal{Y}_\tau^{-1}$ for $\mathcal{Y} \in U(1)^{[\tau]}$. In general, however, they may differ by a 1-cocycle on Z with values in $U(1)^{[\tau]}$. We infer that the cohomology group

$$H^1(Z, U(1)^{[\tau]}) \;\cong\; H^1(Z_{[\tau]}, U(1)) \;=\; Z^*_{[\tau]},$$

where $Z^*_{[\tau]}$ is the character group of $Z_{[\tau]}$, acts freely and transitively on the set of the isomorphism classes of symmetric rank 1 $\mathcal{G}'$-branes supported by $D' = \mathcal{C}'_{[\tau]}$, if this set is non-empty. Recall that $Z_{[\tau]}$ is the fundamental group of the conjugacy class $\mathcal{C}'_{[\tau]} \subset G'$ so that $Z^*_{[\tau]}$ may be viewed as the group of isomorphism classes of flat line bundles over $\mathcal{C}'_{[\tau]}$. Let $\mathcal{E}'_1(h)$, $h = 1, \dots, Z_{[\tau]}$, denote the non-isomorphic 1-dimensional $\mathcal{G}'_{D'}$-modules obtained from non-equivalent solutions $\mathcal{W}$ of Eq. (5.18). They give rise to non-isomorphic rank 1 $\mathcal{G}'$-branes $\mathcal{D}_1(h) = (D', \mathcal{E}'_1(h))$. In such a situation, any N-dimensional $\mathcal{G}'|'_D$-module $\mathcal{E}'_N$ giving rise to a symmetric brane is equal, up to isomorphism, to a product of $\mathcal{E}'_1(h)$ (for any h) by a flat vector bundle over $D' = \mathcal{C}'_{[\tau]}$. Such vector bundles are, again up to isomorphism, direct sums of flat line bundles so we may set

$$\mathcal{E}'_N \;=\; \mathcal{E}'_1(h_1) \oplus \cdots \oplus \mathcal{E}'_1(h_N)\,.$$

We infer that if the cohomology class $[\mathcal{V}]$ is trivial then any rank N $\mathcal{G}'$-brane $\mathcal{D}'_N$ supported by $\mathcal{C}'_{[\tau]}$ is isomorphic to a stack of N (in general, different) 1-dimensional branes $\mathcal{D}'_1(h_n)$, $h_n \in \{1, \dots, Z_{[\tau]}\}$. This is always the situation if $Z_{[\tau]}$ is a cyclic group since then $H^2(Z_{[\tau]}, U(1)) = \{1\}$.

Let us describe more precisely the solutions $\mathcal{W}$ of Eq. (5.18). First note that if two solutions differ by a 2-cocycle $\mathcal{W}'$ that is a coboundary, $\mathcal{W}' = \delta\mathcal{Y}$, then they coincide when restricted to $Z_{[\tau]}$: $\mathcal{W}'_{\tau;z_0} = 1$ (here and below, we use subscript zero for elements of $Z_{[\tau]}$). Conversely, if two solutions coincide if restricted to $Z_{[\tau]}$ then, setting $\mathcal{Y}_{\widehat{\tau}_z} = \mathcal{W}'_{\widehat{\tau},z}$ for a fixed $\widehat{\tau} \in [\tau]$, we have $\mathcal{W}' = \delta\mathcal{Y}$. A solution of Eq. (5.18) satisfies

$$\mathcal{W}_{\widehat{\tau}_z;z_0} \;=\; \mathcal{W}_{\widehat{\tau};zz_0}\,\mathcal{W}^{-1}_{\widehat{\tau};z}\,\mathcal{V}_{\widehat{\tau};z,z_0} \;=\; \mathcal{W}_{\widehat{\tau};z_0}\,\mathcal{V}^{-1}_{\widehat{\tau};z_0,z}\,\mathcal{V}_{\widehat{\tau};z,z_0} \tag{5.19}$$

so that its restriction to $Z_{[\tau]}$ is determined by the solution of the restricted equation with fixed $\widehat{\tau}$

$$(\delta\mathcal{W})_{\widehat{\tau};z_0,z'_0} \;=\; \mathcal{W}_{\widehat{\tau};z'_0}\,\mathcal{W}^{-1}_{\widehat{\tau};z_0z'_0}\,\mathcal{W}_{\widehat{\tau};z_0} \;=\; \mathcal{V}_{\widehat{\tau};z_0,z'_0} \tag{5.20}$$

involving the simpler cohomology of group $Z_{[\tau]}$ with values in $U(1)$. The general solution of the last equation has the form

$$\mathcal{W}_{\widehat{\tau};z_0} \;=\; \mathcal{W}^{\widehat{\tau}}_{\widehat{\tau};z_0}\,\widehat{\chi}(z_0)\,,$$

where $\mathcal{W}^{\widehat{\tau}}_{\widehat{\tau};z_0}$ is a particular solution and $\widehat{\chi}$ is a character of $Z_{[\tau]}$.

Any solution of the restricted equation (5.20) may be extended to a solution of (5.18) in the following way. First, fix a map $[\tau] \ni \tau \longmapsto z(\tau) \in Z$ such that $z(\tau)\tau = \widehat{\tau}$ and, for $\tau \neq \widehat{\tau}$, choose $\mathcal{W}_{\widehat{\tau};z(\tau)} \in U(1)$ arbitrarily. Setting

$$\mathcal{W}_{\widehat{\tau};z_0 z(\tau)} \;=\; \mathcal{W}_{\widehat{\tau};z(\tau)}\,\mathcal{W}_{\widehat{\tau};z_0}\,\mathcal{V}^{-1}_{\widehat{\tau};z_0,z(\tau)}\,,$$

one defines $\mathcal{W}_{\widehat{\tau};z}$ for all $z \in Z$ in such a way that the above equation holds also if $z(\tau)$ is replaced by any $z \in Z$. Finally, with the help of the relation

$$\mathcal{W}_{\widehat{\tau}_z;z'} \;=\; \mathcal{W}_{\widehat{\tau};zz'}\,\mathcal{W}^{-1}_{\widehat{\tau};z}\,\mathcal{V}_{\widehat{\tau};z,z'}$$

one defines a solution $\mathcal{W}$ of Eq. (5.18) which, in particular, satisfies the identity (5.19).

If another fixed element $\widehat{\tau}' = \widehat{\tau}_z$ of the orbit $[\tau]$ is used in the construction above with

$$\mathcal{W}_{\widehat{\tau}',z_0} \;=\; \mathcal{W}^{\widehat{\tau}'}_{\widehat{\tau}',z_0}\,\widehat{\chi}'(z_0)$$

as the solution of the restricted equation then the resulting solutions of the complete equation (5.18) coincides with the one described previously if and only if

$$\widehat{\chi}'(z_0) \;=\; \widehat{\phi}_{\widehat{\tau};z}(z_0)\,\widehat{\chi}(z_0)$$

for

$$\widehat{\phi}_{\widehat{\tau};z}(z_0) \;=\; \mathcal{W}^{\widehat{\tau}}_{\widehat{\tau};z_0}\,{\mathcal{W}^{\widehat{\tau}_z}_{\widehat{\tau}_z;z_0}}^{-1}\,\mathcal{V}_{\widehat{\tau};z,z_0}\,\mathcal{V}^{-1}_{\widehat{\tau};z_0,z}\,, \tag{5.21}$$

as may be easily seen with the help of relation (5.19). Note that $\widehat{\phi}$ has the following properties:

$$\begin{gathered}\widehat{\phi}_{\widehat{\tau};z_0 z}(z_0') = \widehat{\phi}_{\widehat{\tau};z}(z_0')\,, \qquad \widehat{\phi}_{\widehat{\tau};z}(z_0 z_0') \;=\; \widehat{\phi}_{\widehat{\tau};z}(z_0)\,\widehat{\phi}_{\widehat{\tau};z}(z_0')\,,\\ \widehat{\phi}_{\widehat{\tau}_z;z'}(z_0)\,\widehat{\phi}_{\widehat{\tau};zz'}(z_0)^{-1}\,\widehat{\phi}_{\widehat{\tau};z}(z_0) \;=\; 1\,.\end{gathered} \tag{5.22}$$

They mean that $\widehat{\phi}$ defines a 1-cocycle on group Z with values in the module of $Z^*_{[\tau]}$-valued functions on the orbit $[\tau]$ which, due to the first relation, descends to a cocycle on the quotient group $Z/Z_{[\tau]}$. The cocycle $\widehat{\phi}$ is not unique. If we multiply the special solutions $\mathcal{W}^{\widehat{\tau}}_{\widehat{\tau},z_0}$ by $\widehat{\tau}$-dependent characters $\widehat{\chi}_{\widehat{\tau}}(z_0)$ of $\mathcal{Z}_{[\tau]}$ then

$$\widehat{\phi}_{\widehat{\tau};z}(z_0) \;\longmapsto\; \widehat{\phi}_{\widehat{\tau};z}(z_0)\;\widehat{\chi}_{\widehat{\tau}}(z_0)\,\chi_{\widehat{\tau}_z}(z_0)^{-1}\,.$$

Given cocycle $\widehat{\phi}$, we may identify the set of symmetric rank 1 $\mathcal{G}'$-branes supported by $\mathcal{C}'_{[\tau]}$ with the set of equivalence classes $[\widehat{\tau},\widehat{\chi}]$ of pairs $(\widehat{\tau},\widehat{\chi})$ with $\widehat{\tau} \in [\tau]$ and $\widehat{\chi} \in Z^*_{[\tau]}$ such that

$$(\widehat{\tau},\widehat{\chi}) \;\sim\; (\widehat{\tau}',\widehat{\chi}') \quad \text{if} \;\; \widehat{\tau}' = \widehat{\tau}_z \;\; \text{and} \;\; \widehat{\chi}' = \widehat{\phi}_{\widehat{\tau};z}(\,\cdot\,)\,\widehat{\chi}\,.$$

Such a description of branes agrees with the general one conjectured in [30, 66] for conformal field theories of the simple current extension type. The general classification of the branes proposed there, based on consistency considerations, involved equivalence classes of primary fields and characters of their "central stabilizers". For rank 1 branes of the WZW models, the latter reduce to the ordinary stabilizer subgroups $\mathcal{Z}_{[\tau]}$ in the simple current group $\mathcal{Z}$.

In Appendix A we list solutions of Eq. (5.18) giving rise to all (up to isomorphism) symmetric rank 1 $\mathcal{G}'$-branes supported by the conjugacy classes of non-simply connected groups $G' = G/Z$ with covering groups G running through the Cartan list. We use the description of the Weyl alcoves and of choices of the 2-chains b and V entering Eqs. (4.5) and (5.16) taken from [40][6]. We specify:

the action of the center of G on the Weyl alcove,
the restrictions on $n_i \geq 0$, $\sum\limits_{i=0}^{r} n_i = \mathsf{k}$, such that $\sum\limits_{i=0}^{r} n_i\tau_i = \mathsf{k}\tau \equiv \lambda$ is a weight,
the subgroups Z and the restrictions on the level k of gerbes $\mathcal{G}'$ over G/Z,
the 2-chains V,
the 2-cocycles $\mathcal{V}$,
possible stabilizer subgroups of Z,
the special solutions $\mathcal{W}^{\widehat{\tau}}$ of Eq. (5.18) for different Z-orbits $[\tau]$,
general solutions leading to non-isomorphic $\mathcal{G}'$-branes.

Describing 2-cocycle $\mathcal{V}$ we use the the monodromy charge $Q_z(\lambda)$ defined for integrable weights λ by

$$Q_z(\lambda) \;=\; h(\lambda) \,+\, h(k\tau_{z0}) \,-\, h(z\lambda) \;\mathrm{mod}\, 1 \;=\; \mathrm{tr}\,\lambda\tau_{0_z} \;\mathrm{mod}\, 1$$

where $h(\lambda)$ is the conformal weight of the chiral primary field with weight λ and, by definition, $z\lambda = \mathsf{k}z\tau$ if $\lambda = \mathsf{k}\tau$. In most cases, the special solutions $\mathcal{W}^{\widehat{\tau}}_{\tau;z,z'}$ are independent of $\widehat{\tau}, \tau \in [\tau]$ as may be seen from the restrictions on coefficients n_i together with the relation $\sum n_i = \mathsf{k}$ combined with the restrictions on k. Such independence implies that the cocycles $\widehat{\phi}$ of Eq. (5.21) are trivial and the description of branes by the equivalence classes $[\widehat{\tau}, \widehat{\chi}]$ is superfluous. The exception is the group $G' = Spin(4n)/(\mathbb{Z}_2 \times \mathbb{Z}_2) = SO(4n)/\mathbb{Z}_2$ where several cases give rise to non-trivial cocycle $\widehat{\phi}$. Also for that group, which is the only one with non-cyclic fundamental group $Z \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ and multiple (double) choice of stably-non-isomorphic gerbes $\mathcal{G}'$, for one of those gerbes there are no (scalar) solutions of Eq. (5.18) for single-point orbits $[\tau]$ (i.e. if $Z_{[\tau]} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$). Consequently, there are no symmetric rank 1 $\mathcal{G}'$-branes supported by the corresponding (integrable) conjugacy classes $\mathcal{C}'_{[\tau]} \subset G'$. We shall devote the next section to the analysis of higher-rank $\mathcal{G}'$-branes supported by such conjugacy classes.

6 Non-Abelian symmetric branes for $SO(4n)/\mathbb{Z}_2$

For group $G' = Spin(2r)/Z$ with even rank $r = 4, 6, \ldots$ and $Z \cong \mathbb{Z}_2 \times \mathbb{Z}_2$, the integrable conjugacy classes $\mathcal{C}'_{[\tau]}$ with $Z_{[\tau]} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ correspond to 1-point orbits $[\tau]$ with

$$\mathsf{k}\tau \;=\; \sum_{i=0}^{r} n_i\tau_i\,, \qquad\qquad n_i \geq 0\,, \qquad\qquad \sum_{i=0}^{r} n_i \;=\; \mathsf{k}\,,$$

---

[6] $V_{z,z'}$ is equal to $\mathrm{e}^{-2\pi\mathrm{i}\mathsf{k}\,\mathrm{tr}\,\tau_{zz'0}b_{z,z'}}\, u_{z,z'}$ with $u_{z,z'}$ defined and calculated in [40]

$$n_0 = n_1 = n_{r-1} = n_r \in \mathbb{Z}\,, \qquad n_i = n_{r-i} \in 2\mathbb{Z},\ \ i = 2, \dots, r-2\,.$$

Such conjugacy classes exist if and only if k is even. The two inequivalent choices $\mathcal{G}'_\pm$ for the gerbe over G' correspond to the upper and lower choices of signs in the 2-chain V that enters the formula (4.11) for the groupoid multiplication,

$$V_{z,z'} \;=\; \begin{cases} \pm 1 & \text{if} \quad (z,z') = (z_2,z_1), (z_2, z_1z_2), \\ & \qquad\qquad (z_1z_2, z_1), (z_1z_2, z_1z_2)\,, \\ 1 & \text{otherwise} \end{cases} \tag{6.1}$$

where z_1, z_2 are the generators of Z, see Sec. 4.4.II of [40]. For the 1-point orbits $[\tau]$, the 2-cocycle $\mathcal{V}$ of Eq. (5.16) reduces to the same expression

$$\mathcal{V}_{\tau;z,z'} \;=\; V_{z,z} \tag{6.2}$$

and defines for the lower choice of signs in 6.1 a non-trivial cohomology class $[\mathcal{V}] \in H^2(\mathbb{Z}_2 \times \mathbb{Z}_2, U(1)) \cong \mathbb{Z}_2$ which is an obstruction to the existence of solutions of Eq. (5.18) and of symmetric rank 1 $\mathcal{G}'_-$-branes supported by the conjugacy class $\mathcal{C}'_{[\tau]}$. The simplest instances with $\mathsf{k} > 0$ occur for group $G' = Spin(8)/(\mathbb{Z}_2 \times \mathbb{Z}_2) = SO(8)/\mathbb{Z}_2$ for $\tau = \tau_2$ at levels $\mathsf{k} = 2, 4, \dots$ and for $\tau = \frac{1}{4}(\tau_0 + \tau_1 + \tau_3 + \tau_4)$ at $\mathsf{k} = 4, 6, \dots$ in conventions of [40].

We shall search now for symmetric rank N $\mathcal{G}'_-$-branes supported by the integrable conjugacy classes $\mathcal{C}'_{[\tau]}$ with 1-point orbits $[\tau]$. As in the rank 1 case, such a brane $\mathcal{D}'_N = (D', \mathcal{E}'_N)$ induces by restriction a brane $(D, \mathcal{E}_N)$ for the $Spin(2r)$ theory with the support D given by the conjugacy class $\mathcal{C}_\tau \subset Spin(2r)$. If $\mathcal{E}'_N = (E', \rho')$ then $\mathcal{E}_N = (E, \rho)$ with coinciding vector bundles $E' = E$ over $Y'_{D'} = Y_D$. Since the symmetric rank N $\mathcal{G}$-branes are isomorphic to stacks of N rank 1 branes, we may assume, passing at most to an isomorphic $\mathcal{G}'_-|_{D'}$-module $\mathcal{E}'_N$, that over each connected component $P_i|_{\mathcal{C}_\tau} \cong Y_{i\tau}$ of $Y'_{D'}$, see (5.8), the vector bundle E' restricts to the bundle $L_{i\tau} \otimes \mathbb{C}^N$. Elements $\ell_{i\tau} \in L_{i\tau} \otimes \mathbb{C}^N$ in the fiber over $y = (g, h\gamma^{-1}) \in P_i|_{\mathcal{C}_\tau}$ are now determined by the equivalence classes $\widehat{\ell}_{i\tau} = [\gamma, v]_i$ defined as in (5.5) but with $v \in \mathbb{C}^N$ rather than $v \in \mathbb{C}$. With this modification, the discussion around Eqs. (5.9) to (5.16) may be repeated word by word with $\mathcal{W}^{ij}_{\tau;z}$ and $\mathcal{Y}^i_\tau$ being now unitary $N \times N$ matrices so that their order in the formulae, previously irrelevant, becomes important. The same argument as before shows that $\mathcal{W}^{ij}_{\tau;z}$ may be chosen (i,j)-independent. Eq. (5.15) reduces then to the matrix relation

$$\mathcal{W}_{\tau;z}\, \mathcal{W}_{\tau;z'} \;=\; \mathcal{V}_{\tau;z,z'}\, \mathcal{W}_{\tau;zz'} \tag{6.3}$$

which, with the use of Eqs. (6.2) and (6.1) with the lower choice for the signs, implies that $\mathcal{W}_{\tau;1} = 1$ and that

$$\mathcal{W}^2_{\tau;z_1} \;=\; 1 \;=\; \mathcal{W}^2_{\tau;z_2}\,, \qquad \mathcal{W}_{\tau;z_1}\, \mathcal{W}_{\tau;z_2} + \mathcal{W}_{\tau;z_2}\, \mathcal{W}_{\tau;z_1} \;=\; 0$$

with $\mathcal{W}_{\tau;z_1z_2} = \mathcal{W}_{\tau;z_1} \mathcal{W}_{\tau;z_2}$. In other words, matrices $\mathcal{W}_{\tau;z}$ define an N-dimensional representation of the 2-dimensional Clifford algebra. Unitarily equivalent representations correspond to isomorphic $\mathcal{G}'_-|_{D'}$ modules i.e. to isomorphic $\mathcal{G}'_-$-branes and *vice versa*. The lowest dimensional representation is of dimension 2 and is given by the Pauli matrices

$$\mathcal{W}_{\tau;\,z_1} \;=\; \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \qquad \mathcal{W}_{\tau;z_2} \;=\; \begin{pmatrix} 0 & -\mathsf{i} \\ \mathsf{i} & 0 \end{pmatrix}, \qquad \mathcal{W}_{\tau;z_1z_2} \;=\; \mathsf{i} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \tag{6.4}$$

It is unique up to unitary equivalence. It gives rise to a 2-dimensional $\mathcal{G}'_-|_{D'}$-module $\mathcal{E}'_2$, hence to a rank 2 $\mathcal{G}'_-$-brane $\mathcal{D}'_2$. The higher dimensional representations exist in even dimensions N and are unitarily equivalent to direct sums of the 2-dimensional representation so that, up to isomorphism,

$$\mathcal{E}'_N \;=\; \underbrace{\mathcal{E}'_2 \oplus \cdots \oplus \mathcal{E}'_2}_{N/2 \text{ times}} \,.$$

We infer that for the gerbe $\mathcal{G}'_-$ over $SO(4n)/\mathbb{Z}_2$ corresponding to the lower choice of signs in (6.1) and even level k, the symmetric rank N $\mathcal{G}'_-$-branes supported by the integrable conjugacy classes $\mathcal{C}'_{[\tau]}$ with 1-point orbits $[\tau]$ exist only for N even and are isomorphic to a stack of $N/2$ rank 2 non-Abelian $\mathcal{G}'_-$-branes $\mathcal{D}'_2$ determined by the solution (6.4) of Eq. (6.3). In the algebraic classification of branes [30, 66, 70], see also [65], based on the simple current technique, those are the cases where the "central stabilizer" subgroup of the simple current group Z is trivial hence smaller than the ordinary stabilizer $Z_{[\tau]}$. That the branes corresponding to such situations should be counted among the boundary conditions follows by demanding that the total number of such conditions be equal to the dimension I of the set of the Ishibashi states in the bulk sector of the theory [30, 66]. The simplest example of such a count is the case of group $G' = SO(8)/\mathbb{Z}_2$ at level $\mathsf{k} = 2$ where the space of Ishibashi states has dimension $I_+ = 11$ and $I_- = 8$ for the bulk theories related to gerbes $\mathcal{G}'_\pm$, respectively. There are five orbits $[\tau]$ in the Weyl alcove $\mathcal{A}_W$ such that $\mathsf{k}\tau$ is a weight. With the choice of $\mathcal{A}_W$ and its vertices τ_i as in [40] leading to the Z-action $z_1\tau_i = \tau_{4-i}$ $z_2\tau_0 = \tau_1$, $z_2\tau_2 = \tau_2$, $z_2\tau_3 = \tau_4$, these orbits are:

$$\begin{gathered}\{\tau_0,\,\tau_1,\,\tau_3,\,\tau_4\},\\ \{\tfrac{1}{2}\tau_0+\tfrac{1}{2}\tau_1,\,\tfrac{1}{2}\tau_3+\tfrac{1}{2}\tau_4\},\quad \{\tfrac{1}{2}\tau_0+\tfrac{1}{2}\tau_3,\,\tfrac{1}{2}\tau_1+\tfrac{1}{2}\tau_4\},\quad \{\tfrac{1}{2}\tau_0+\tfrac{1}{2}\tau_4,\,\tfrac{1}{2}\tau_1+\tfrac{1}{2}\tau_3\},\\ \{\tau_2\}\,.\end{gathered}$$

For the boundary theory related to gerbe $\mathcal{G}'_+$, each of those corresponds to integrable conjugacy class carrying $|Z_{[\tau]}|$ non-isomorphic symmetric rank 1 branes, so altogether we obtain $1+2+2+2+4 = 11$ branes. For the theory related to $\mathcal{G}'_-$, the conjugacy class corresponding to the 1-point orbit $\{\tau_2\}$ gives rise to a rank 2 brane unique up to isomorphism and the brane count becomes $1+2+2+2+1 = 8$.

7 Boundary partition functions and operator product

7.1 Boundary states for simply connected target groups

Although obtained by classical considerations, the geometric classification of branes in the WZW model permits to elucidate the quantized theory by providing structures that manifest themselves directly on the quantum level. The spaces of states in the quantum WZW theory may be realized with the use of geometric quantization as spaces of sections of vector bundles over the group path spaces. Such vector bundles are canonically associated to the gerbe $\mathcal{G}$ on the group and to pairs of $\mathcal{G}$-branes $\mathcal{D}^s$, $s = 0, 1$. More concretely, for $\mathcal{L}$ the line bundle over the loop group LG and $\mathcal{E}^{\mathcal{D}^1}_{\mathcal{D}^0}$ the vector bundle over the space of open paths $IG^{D^1}_{D^0}$ in the group extending between the brane supports, see Sect. 3.2, the spaces of sections[7]

$$\mathbb{H}\ =\ \Gamma(\mathcal{L})\,,\qquad\quad \mathbb{H}^{\mathcal{D}^1}_{\mathcal{D}^0}\ =\ \Gamma(\mathcal{E}^{\mathcal{D}^1}_{\mathcal{D}^0})\tag{7.1}$$

provide, respectively, the spaces of bulk and of boundary states of the theory. The first one carries the action of a left-right current algebra $\widehat{\mathbf{g}}\oplus\widehat{\mathbf{g}}$ whereas the second one that of the chiral current algebra $\widehat{\mathbf{g}}$. The analysis of the representation content (spectrum) of the space of bulk states for the WZW models with all compact, simple but possibly non-simply connected groups has been performed in [27]. For the space of boundary states, the decomposition into the irreducible highest weight representations of the current algebras has the form

$$\mathbb{H}^{\mathcal{D}^1}_{\mathcal{D}^0}\ \cong\ \mathop{\oplus}_{\lambda}\Big(M^{\mathcal{D}^1}_{\mathcal{D}^0\lambda}\otimes\widehat{V}_{\bar\lambda}\Big)\tag{7.2}$$

where the direct sum is over the integrable weights λ and spaces $\widehat{V}_{\bar\lambda}$ carry the irreducible representations of the current algebra $\widehat{\mathbf{g}}$ of level k and highest weight $\bar\lambda$[8]. For the theory with the simply connected target group G and rank N_s symmetric $\mathcal{G}$-branes $\mathcal{D}^s$ that are stacks of rank 1 branes supported by the integrable conjugacy classes C_{τ^s} corresponding to weights $\lambda^s=\mathsf{k}\tau^s$, the multiplicity spaces take the product form

$$M^{\mathcal{D}^1}_{\mathcal{D}^0\lambda}\ =\ \mathbb{C}^{N_0}\otimes\overline{\mathbb{C}^{N_1}}\otimes M^{\lambda^1}_{\lambda^0\lambda}\,.\tag{7.3}$$

The dimensions of the multiplicity spaces $M^{\lambda^1}_{\lambda^0\lambda}$ for the rank 1 branes are equal to the fusion coefficients $N^{\lambda^1}_{\lambda^0\lambda}$ so that the **boundary partition functions** of the WZW theory with a simply connected target group are given by the expressions

$$Z^{\mathcal{D}^1}_{\mathcal{D}^0}(T)\ \equiv\ \mathrm{Tr}_{\mathbb{H}^{\mathcal{D}^1}_{\mathcal{D}^0}}\,\mathrm{e}^{-TL_0}\ =\ N_0N_1\sum_{\lambda}N^{\lambda^1}_{\lambda^0\lambda}\,\widehat{\chi}_{\bar\lambda}(\tfrac{\mathrm{i}T}{2\pi})$$

where the (restricted) affine characters $\widehat{\chi}_{\lambda}(\frac{\mathrm{i}T}{2\pi})\ =\ \mathrm{Tr}_{\widehat{V}_\lambda}\mathrm{e}^{-TL_0}$ with L_0 denoting the Virasoro generator given by the Sugawara construction.

The multiplicity spaces $M^{\lambda^1}_{\lambda^0\lambda}$ may be thought of in different ways, depending on the situation. Firstly, they may be naturally identified [41, 38] with the spaces of 3-point genus zero conformal blocks of the bulk, group G WZW theory with insertions of the primary fields corresponding to the integrable weights λ^1, $\bar\lambda^0$ and $\bar\lambda$ or, in other words, as spaces of properly defined intertwiners of the current algebra $\widehat{\mathbf{g}}$ actions in the spaces $\widehat{V}_{\lambda^1}$ and $\widehat{V}_{\lambda^0}\otimes\widehat{V}_{\lambda}$. Secondly, they may be identified with the "fusion rule" subspace of the intertwiners of the Lie algebra $\mathbf{g}$ action

$$\mathrm{Hom}^{\mathrm{FR}}_{\mathbf{g}}(V_{\lambda^1},V_{\lambda^0}\otimes V_\lambda)\ \subset\ \mathrm{Hom}_{\mathbf{g}}(V_{\lambda^1},V_{\lambda^0}\otimes V_\lambda)$$

where V_λ stands for the space of the irreducible highest weight λ representation of $\mathbf{g}$. The relation with the previous picture is that the Lie algebra intertwiners in the fusion rule subspace are the ones that extend to the current algebra intertwiners (such extensions are unique). Finally, the spaces $M^{\lambda^1}_{\lambda^0\lambda}$ may be thought of as composed of the

---

[7] For positivity of energy, the level k of the theory has to be taken positive

[8] $\bar\lambda$ denotes the highest weight of the irreducible representation of $\mathbf{g}$ complex conjugate to the one with the highest weight λ.

("good") intertwiners of the action of the quantum deformation $\mathcal{U}_q(\mathbf{g})$ of the enveloping algebra of $\mathbf{g}$ in its highest weight modules.

Different realizations of spaces $M^{\lambda^1}_{\lambda^0\lambda}$ are more convenient in different contexts. We shall need still another realization that is derived from the one based on the fusion rule intertwiners of the Lie algebra action. We shall need a more concrete description of the spaces $\mathrm{Hom}^{\mathrm{FR}}_{\mathbf{g}}(V_{\lambda^1}, V_{\lambda^0}\otimes V_\lambda)$. Consider the linear mapping

$$\mathrm{Hom}_{\mathbf{g}}(V_{\lambda^1}, V_{\lambda^0}\otimes V_\lambda)\ \ni\ \psi\ \longmapsto\ |\psi\rangle\ \in\ V_\lambda$$

such that for all $|v\rangle\in V_\lambda$ and the highest weight vectors $|\lambda^s\rangle\in V_{\lambda^s}$ annihilated by the step generators e_α for positive roots α of $\mathbf{g}$,

$$\langle v\,|\,\psi\rangle\ =\langle\lambda^0\otimes v\,|\,\psi\,|\lambda^1\rangle\,.$$

It is easy to see that $|\psi\rangle$ determines the intertwiner ψ uniquely so that $\mathrm{Hom}_{\mathbf{g}}(V_{\lambda^1}, V_{\lambda^0}\otimes V_\lambda)$ and its fusion rule subspace may be identified with subspaces in V_λ. The latter is characterized by the conditions

$$t\,|\psi\rangle\ =\ \mathrm{tr}\,t(\lambda^1-\lambda^0)\,|\psi\rangle\qquad\text{for}\quad t\in\mathbf{t}\,,\tag{7.4}$$

$$e_{-\alpha_i}^{\mathrm{tr}\,\alpha_i^\vee\lambda^1+1}\,|\psi\rangle\ =\ 0\qquad\text{for}\quad i=1,\dots,r\,,\tag{7.5}$$

$$e_{\phi}^{\mathsf{k}-\mathrm{tr}\,\phi^\vee\lambda^1+1}\,|\psi\rangle\ =\ 0\tag{7.6}$$

whereas for the former, the 3$^{\text{rd}}$ condition should be dropped. Above $\mathbf{t}$ denotes the Cartan subalgebra of $\mathbf{g}$, $\alpha_i^\vee$ the simple coroots and ϕ, $\phi^\vee$ the highest root of $\mathbf{g}$ and its coroot. Equivalently, one may replace conditions (7.5) and (7.6) by

$$\begin{aligned}e_{\alpha_i}^{\mathrm{tr}\,\alpha_i^\vee\lambda^0+1}\,|\psi\rangle\ &=\ 0\qquad\text{for}\quad i=1,\dots,r\,,\\ e_{-\phi}^{\mathsf{k}-\mathrm{tr}\,\phi^\vee\lambda^0+1}\,|\psi\rangle\ &=\ 0\,.\end{aligned}$$

Below, we shall identify the multiplicity space $M^{\lambda^1}_{\lambda^0\lambda}$ with the subspace of V_λ composed of vectors satisfying (7.4), (7.5) and (7.6):

$$M^{\lambda^1}_{\lambda^0\lambda}\ \subset\ V_\lambda\,.\tag{7.7}$$

$M^{\lambda^1}_{\lambda^0\lambda}$ inherits this way the scalar product from V_λ.

The latter description of the multiplicity spaces is particularly natural in the geometric realization (7.1) of the spaces of states where it may be obtained by evaluating sections of the vector bundle $\mathcal{E}^{\mathcal{D}^1}_{\mathcal{D}^0}$ on the special path

$$[0,\pi]\ \ni\ x\ \mapsto\ g_{\tau^0\tau^1}(x)=\mathrm{e}^{2\pi\mathrm{i}(\tau^0+x(\tau^1-\tau^0)/\pi)}$$

in G extending between the brane supports D^0 and D^1. Such evaluation takes values in the fiber of $\mathcal{E}^{\mathcal{D}^1}_{\mathcal{D}^0}$ over $g_{\tau^0\tau^1}$ which may be canonically identified with $\mathbb{C}^{N_0}\otimes\overline{\mathbb{C}^{N_1}}$. The composition of natural injections

$$\begin{aligned}\mathrm{Hom}_{\widehat{\mathbf{g}}}\Big(\widehat{V}_{\bar\lambda},\,\Gamma(\mathcal{E}^{\mathcal{D}^1}_{\mathcal{D}^0})\Big)\ &\hookrightarrow\ \mathrm{Hom}_{\mathbf{g}}\Big(V_{\bar\lambda},\,\Gamma(\mathcal{E}^{\mathcal{D}^1}_{\mathcal{D}^0})\Big)\ \hookrightarrow\ \Gamma(\mathcal{E}^{\mathcal{D}^1}_{\mathcal{D}^0})\otimes V_\lambda\\ &\hookrightarrow\ \mathbb{C}^{N_0}\otimes\overline{\mathbb{C}^{N_1}}\otimes V_\lambda\,,\end{aligned}\tag{7.8}$$

where the last but one injection uses the identification between the dual space of $V_{\bar\lambda}$ and V_λ and the last one evaluates the sections of $\mathcal{E}^{\mathcal{D}^1}_{\mathcal{D}^0}$ at $g_{\tau^0\tau^1}$, embeds the multiplicity space into $\overline{\mathbb{C}^{N_1}}\otimes\mathbb{C}^{N_0}\otimes V_\lambda$.

7.2 Boundary states for non-simply connected target groups

As pointed out in Sect. 9.2 of [39], the geometric realization (7.1) of the boundary spaces of states implies a simple relation between those spaces for the WZW models with a simply-connected target group G and with the quotient target $G' = G/Z$. In one phrase, the latter space of states is composed of the Z-invariant states of the former. More exactly, for symmetric $\mathcal{G}'$-branes $\mathcal{D}'^s$, $s = 0, 1$, supported by the integrable conjugacy classes $\mathcal{C}'_{[\tau^s]} \subset G'$, let $\mathcal{D}^s$ run through the $\mathcal{G}$-branes supported by the conjugacy classes $\mathcal{C}_{\tau^s}$ with $\tau^s \in [\tau^s]$ obtained by the pullback of branes $\mathcal{D}'^s$. There is one such $\mathcal{G}$-brane $\mathcal{D}^s$ for each $\tau^s \in [\tau^s]$. We shall denote by $\mathcal{D}^s_z$ the brane supported by $z^{-1}\mathcal{C}_{\tau^s} = \mathcal{C}_{\tau^s_z}$ if $\mathcal{D}^s$ is supported by τ^s. As has been discussed in [39], the $\mathcal{D}'^s$ brane structures allow to lift naturally the action of the subgroup Z of the center of G to the vector bundles $\mathcal{E}^{\mathcal{D}^1}_{\mathcal{D}^0}$:

$$\begin{tikzcd}
\mathcal{E}^{\mathcal{D}^1_z}_{\mathcal{D}^0_z} \arrow[r, "\widehat{z}"] \arrow[d] & \mathcal{E}^{\mathcal{D}^1}_{\mathcal{D}^0} \arrow[d] \\
IG^{D^1_z}_{D^0_z} \arrow[r, "z"] & IG^{D^1}_{D^0}
\end{tikzcd}$$

with the lower line given by multiplication by z. The action of Z on the vector bundles $\mathcal{E}^{\mathcal{D}^1}_{\mathcal{D}^0}$ satisfies $\widehat{z}\widehat{z}' = \widehat{zz'}$ and induces the action on the sections:

$$(\mathbb{U}(z)\Psi)(g) \ = \ \widehat{z}\,\Psi(z^{-1}g)$$

for $\Psi \in \Gamma(\mathcal{E}^{\mathcal{D}^1_z}_{\mathcal{D}^0_z})$. One obtains in this way a representation $\mathbb{U}$ of Z in the space

$$\widetilde{\mathbb{H}}'^{\mathcal{D}'^1}_{\mathcal{D}'^0} \ = \ \underset{(\mathcal{D}^0,\mathcal{D}^1)}{\oplus} \mathbb{H}^{\mathcal{D}^1}_{\mathcal{D}^0}$$

containing all the G-theory states compatible with the ones in $\mathbb{H}'^{\mathcal{D}'^1}_{\mathcal{D}'^0}$. Operators $\mathbb{U}(z)$ map $\mathbb{H}^{\mathcal{D}^1_z}_{\mathcal{D}^0_z}$ to $\mathbb{H}^{\mathcal{D}^1}_{\mathcal{D}^0}$. They commute with the action of the current algebra so that through the decomposition (7.2) they induces the maps

$$U^{\mathcal{D}^1}_{\mathcal{D}^0\lambda}(z) \ : \ \mathbb{C}^{N_0} \otimes \overline{\mathbb{C}^{N_1}} \otimes M^{\lambda^1_z}_{\lambda^0_z\lambda} \ \longrightarrow \ \mathbb{C}^{N_0} \otimes \overline{\mathbb{C}^{N_1}} \otimes M^{\lambda^1}_{\lambda^0\lambda}$$

on the multiplicity spaces where $z\lambda = \mathsf{k}z\tau$ for $\lambda = \mathsf{k}\tau$ and $\lambda_z = \mathsf{k}\tau_z = \mathsf{k}z^{-1}\tau$. A closer inspection of the the embeddings (7.8) shows that, in the realization (7.7) of spaces $M^{\lambda^1}_{\lambda^0\lambda}$,

$$U^{\mathcal{D}^1}_{\mathcal{D}^0\lambda}(z) \ = \ \mathcal{W}^0_{\lambda^0;z} \ \otimes \ \overline{\mathcal{W}^1_{\lambda^1;z}} \ \otimes \ w_z \tag{7.9}$$

where $\mathcal{W}^s_{\lambda^s;z}$ are the solutions of the cohomological equation (5.18) (with values in $N_s \times N_s$ matrices) relabeled, for notational convenience, by the weights $\lambda^s = \mathsf{k}\tau^s$. Recall that the solutions $\mathcal{W}^s$ define the $\mathcal{G}'|_{\mathcal{D}'^s}$-modules $\mathcal{E}'^s$ such that $\mathcal{D}'^s = (\mathcal{C}'_{[\tau^s]}, \mathcal{E}'^s)$. The elements $w_z \in G$ are as in (4.4). Their action in V_λ intertwines the multiplicity spaces:

$$w_z \ : \ M^{\lambda^1_z}_{\lambda^0_z\lambda} \ \longrightarrow \ M^{\lambda^1}_{\lambda^0\lambda}\,. \tag{7.10}$$

A direct check of this fact may be found in Appendix B. The action (7.10) is a special case of the action of outer current-algebra automorphisms on spaces of conformal blocks studied in [33], see also [1].

The composition rule that assures that $\mathbb{U}$ is a representation of Z,

$$U^{\mathcal{D}^1}_{\mathcal{D}^0\lambda}(z)\; U^{\mathcal{D}^1_z}_{\mathcal{D}^0_z\lambda}(z') \;=\; U^{\mathcal{D}^1}_{\mathcal{D}^0\lambda}(zz')\,,$$

follows easily from the relation $w_z w'_z = \mathrm{e}^{2\pi\mathrm{i}\, b_{z.z'}} w_{zz'}$ and the fact that $\mathrm{e}^{2\pi\mathrm{i}\, b_{z,z'}}$ acts on on $M^{\lambda^1}_{\lambda^0\lambda} \subset V_\lambda$ as the multiplication by $\mathrm{e}^{2\pi\mathrm{i}\,\mathrm{tr}\,(\lambda_1-\lambda_0)b_{z,z'}} = \mathcal{V}_{\lambda^0;z,z'}\,\overline{\mathcal{V}_{\lambda^1;z,z'}}$, see Eqs. (7.4) and (5.16). Operators (7.9) are unitary. They behave naturally under the changes of the solutions $\mathcal{W}^s$ by coboundaries. Such changes lead to isomorphic modules $\mathcal{E}'^s$ and to isomorphisms of vector bundles $\mathcal{E}^{\mathcal{D}^1}_{\mathcal{D}^0}$ and of multiplicity spaces $M^{\mathcal{D}^1}_{\mathcal{D}^0\lambda}$. Operators $\mathbb{U}(z)$ and $U^{\mathcal{D}^1}_{\mathcal{D}^0\lambda}(z)$ change by conjugation with unitary operators induced by such isomorphisms. Similarly, a change $w_z \mapsto \mathrm{e}^{2\pi\mathrm{i}a_z} w_z$ for $a_z \in \mathbf{t}$ induces the change $\mathcal{W}_{\lambda;z} \mapsto \mathrm{e}^{2\pi\mathrm{i}\,\mathrm{tr}\,\lambda a_z}\mathcal{W}_{\lambda;z}$, see Eq. (5.17), that compensates that of w_z in Eq. (7.9).

As indicated in [39], the space of boundary states $\mathbb{H}'^{\mathcal{D}'^1}_{\mathcal{D}'^0}$ of the WZW model with the G' target may be identified with the Z-invariant subspace of $\widetilde{\mathbb{H}}'^{\mathcal{D}'^1}_{\mathcal{D}'^0}$:

$$\mathbb{H}'^{\mathcal{D}'^1}_{\mathcal{D}'^0} \;=\; \tfrac{1}{|Z|}\sum\limits_{z\in Z}\mathbb{U}(z)\;\widetilde{\mathbb{H}}'^{\mathcal{D}'^1}_{\mathcal{D}'^0} \;\equiv\; \mathbb{P}\;\widetilde{\mathbb{H}}'^{\mathcal{D}'^1}_{\mathcal{D}'^0} \tag{7.11}$$

where $\mathbb{P}$ is the orthogonal projector on the Z-invariant states. The scalar product in $\mathbb{H}'^{\mathcal{D}'^1}_{\mathcal{D}'^0}$ should be divided by $|Z|$ with respect to the one inherited from the subspace of $\widetilde{\mathbb{H}}'^{\mathcal{D}'^1}_{\mathcal{D}'^0}$. In terms of the decomposition into the highest weight representations of the current algebra $\widehat{\mathbf{g}}$,

$$\widetilde{\mathbb{H}}'^{\mathcal{D}'^1}_{\mathcal{D}'^0} \;=\; \widetilde{M}'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda}\otimes\widehat{V}_{\bar\lambda} \tag{7.12}$$

with the multiplicity spaces

$$\widetilde{M}'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda} \;=\; \mathop{\oplus}\limits_{(\mathcal{D}^0,\mathcal{D}^1)} M^{\mathcal{D}^1}_{\mathcal{D}^0\lambda} \;=\; \mathop{\oplus}\limits_{\substack{(\lambda^0,\lambda^1)\\ \lambda^s\in[\lambda^s]}} \mathbb{C}^{N_0}\otimes\overline{\mathbb{C}^{N_1}}\otimes M^{\lambda^1}_{\lambda^0\lambda} \tag{7.13}$$

where $[\lambda]$ denotes the Z-orbit of weight λ under the action $\lambda\mapsto z\lambda$. The representation $\mathbb{U}$ induces the operators

$$U'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda}(z) \;=\; \mathop{\oplus}\limits_{(\mathcal{D}^0,\mathcal{D}^1)} U^{\mathcal{D}^1}_{\mathcal{D}^0\lambda}(z) \tag{7.14}$$

providing a unitary representations of Z in the multiplicity spaces $\widetilde{M}'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda}$. The relation (7.11) and the decomposition (7.12) imply then that

$$\mathbb{H}'^{\mathcal{D}'^1}_{\mathcal{D}'^0} \;=\; M'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda}\otimes\widehat{V}_{\bar\lambda}$$

where

$$M'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda} \;=\; \tfrac{1}{|Z|}\sum\limits_{z\in Z} U'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda}(z)\;\widetilde{M}'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda} \;\equiv\; P_\lambda\,\widetilde{M}'^{\mathcal{D}^1}_{\mathcal{D}^0\lambda}\,. \tag{7.15}$$

In short, the multiplicity spaces for the group G' theory are the Z-invariant subspaces of the direct sum of the multiplicity spaces for all group G branes compatible with those of the G' theory.

Together with the representation (7.9), the above constructions lead to the following expression for the boundary partition functions of the group G' WZW theory:

$$Z'^{\mathcal{D}'_1}_{\mathcal{D}'_0}(T) \;\equiv\; \mathrm{Tr}_{\mathbb{H}'^{\mathcal{D}'_1}_{\mathcal{D}'^0}}\, \mathrm{e}^{-T L_0} \;=\; \sum_\lambda N'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda}\; \widehat\chi_{\bar\lambda}(\tfrac{\mathrm{i}T}{2\pi})$$

where

$$\begin{aligned} N'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda} \;&=\; \dim\Big(M'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda}\Big) \;=\; \frac{1}{|Z|}\sum_{z\in Z} \mathrm{Tr}\; U'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda}(z) \\ &=\; \frac{1}{|Z|}\sum_{\lambda^s\in[\lambda^s]}\;\sum_{z\in Z_{[\lambda^0]}\cap Z_{[\lambda^1]}}\;\sum_\lambda\; \mathrm{Tr}\,\mathcal{W}_{\lambda^0;z}\;\; \mathrm{Tr}\,\overline{\mathcal{W}_{\lambda^1;z}}\;\; \mathrm{Tr}_{M^{\lambda^1}_{\lambda^0\lambda}}\, w_z\,. \end{aligned}$$

The non-negative integers $N'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda}$ are the entries of matrices N'_λ that provide a "NIM **representation**" of the fusion algebra [7, 59]. The relation of the above formulae to those of [30, 66] was discussed in [39]. It is based on the Verlinde type expressions conjectured in [33] for the traces of the action of outer automorphisms on the spaces of conformal blocks, see Eq. 9.62 in [39]. It should be possible to provide a direct proof of such trace formulae by studying the action of w_z on $M^{\lambda}_{\lambda^0\lambda}\subset V_\lambda$ (or on the quantum group $\mathcal{U}_q(\mathbf{g})$ realization thereof).

7.3 Boundary operator product

As for the boundary operator product for the WZW theory, it is provided by a linear mapping

$$\mathbb{A}^{\mathcal{D}^1\mathcal{D}^2}_{\mathcal{D}^0\mathcal{D}^1} \,:\, \mathbb{H}^{\mathcal{D}_1}_{\mathcal{D}^0}\otimes\mathbb{H}^{\mathcal{D}_2}_{\mathcal{D}^1} \;\longrightarrow\; \mathbb{H}^{\mathcal{D}_2}_{\mathcal{D}^0}\,.$$

In terms of the multiplicity spaces of the decomposition (7.2), it is determined by the maps

$$A^{\mathcal{D}^1\mathcal{D}^2\,\nu}_{\mathcal{D}^0\mathcal{D}^1\lambda\mu} \,:\, M^{\mathcal{D}^1}_{\mathcal{D}^0\lambda}\otimes M^{\mathcal{D}^2}_{\mathcal{D}^1\mu} \;\longrightarrow\; M^{\mathcal{D}^2}_{\mathcal{D}^0\nu}\otimes \mathrm{Hom}^{\mathrm{FR}}_{\mathbf{g}}(V_\nu, V_\lambda\otimes V_\mu) \tag{7.16}$$

which upon a choice of basis become "**operator product coefficients**". The last factor in (7.16) determines the current-algebra intertwiner from $\widehat{V}_{\bar\lambda}\otimes\widehat{V}_{\bar\mu}$ to $\widehat{V}_{\bar\nu}$ and plays in what follows a spectator role. For simply connected target groups G and rank 1 branes supported by integral conjugacy classes $\mathcal{C}_{\tau^s}$ with $\lambda^s=\mathsf{k}\tau^s$, the maps (7.16) reduce to

$$A^{\lambda^1\lambda^2\,\nu}_{\lambda^0\lambda^1\lambda\,\mu} \,:\, M^{\lambda^1}_{\lambda^0\lambda}\otimes M^{\lambda^2}_{\lambda^1\mu} \;\longrightarrow\; M^{\lambda^2}_{\lambda^0\nu}\otimes M^{\nu}_{\lambda\mu} \tag{7.17}$$

or, in the realization of the latter spaces as fusion rule Lie algebra intertwiners, to the maps

$$\begin{aligned} F_{\lambda^1\nu}\!\begin{bmatrix}\lambda^0 & \lambda^2\\ \lambda & \mu\end{bmatrix} \;:\;\; &\mathrm{Hom}^{\mathrm{FR}}_{\mathbf{g}}(V_{\lambda^1}, V_{\lambda^0}\otimes V_\lambda)\,\otimes\,\mathrm{Hom}^{\mathrm{FR}}_{\mathbf{g}}(V_{\lambda^2}, V_{\lambda^1}\otimes V_\mu) \\ \longrightarrow\;\;\; &\mathrm{Hom}^{\mathrm{FR}}_{\mathbf{g}}(V_{\lambda^2}, V_{\lambda^0}\otimes V_\nu)\,\otimes\,\mathrm{Hom}^{\mathrm{FR}}_{\mathbf{g}}(V_\nu, V_\lambda\otimes V_\mu) \end{aligned}$$

that may be identified with the fusing F-matrices of the bulk WZW theory [55, 63, 25]. Finally, in the realization of the spaces $M^{\nu}_{\lambda\mu}$ as intertwiners of the quantum group $\mathcal{U}_q(\mathbf{g})$

action, the maps (7.17) become [4] the quantum "$6j$ symbols" [18] identifying two ways to decompose triple tensor products of highest weight representations.

For stacks of rank 1 branes in the model with a simply connected target, the maps (7.16) for the multiplicity spaces given by (7.3) are obtained from those in (7.17) by tensoring with the contraction

$$J\,:\,\mathbb{C}^{N_0}\otimes\overline{\mathbb{C}^{N_1}}\otimes\mathbb{C}^{N_1}\otimes\overline{\mathbb{C}^{N_2}}\ \longrightarrow\ \mathbb{C}^{N_0}\otimes\overline{\mathbb{C}^{N_2}}$$

of the middle factors employing the standard scalar product on $\mathbb{C}^{N_1}$.

As pointed out in [39], the picture of the boundary states of the WZW model with the $G'=G/Z$ target as Z-invariant states in the G target model leads to a simple relation between the boundary operator products in both theories. In short, one should project the operator product of the G model to the Z-invariant sector. More precisely, consider the space

$$\widetilde{\mathbb{H}}'^{\mathcal{D}'^1\mathcal{D}'^2}_{\mathcal{D}'^0\mathcal{D}'^1}\ =\ \mathop{\oplus}\limits_{(\mathcal{D}^0,\mathcal{D}^1,\mathcal{D}^2)}\mathbb{H}^{\mathcal{D}^1}_{\mathcal{D}^0}\otimes\mathbb{H}^{\mathcal{D}^2}_{\mathcal{D}^1}\ \subset\ \widetilde{\mathbb{H}}'^{\mathcal{D}'^1}_{\mathcal{D}'^0}\otimes\widetilde{\mathbb{H}}'^{\mathcal{D}'^2}_{\mathcal{D}'^1}$$

where the direct sum is over the pullback branes in the G model and let

$$\widetilde{\mathbb{P}}\,:\,\widetilde{\mathbb{H}}'^{\mathcal{D}'^1}_{\mathcal{D}'^0}\otimes\widetilde{\mathbb{H}}'^{\mathcal{D}'^2}_{\mathcal{D}'^1}\ \longrightarrow\ \widetilde{\mathbb{H}}'^{\mathcal{D}'^1\mathcal{D}'^2}_{\mathcal{D}'^0\mathcal{D}'^1}$$

be the orthogonal projection. Then the boundary operator product of the G' model is given by

$$\mathbb{A}'^{\mathcal{D}'^1\mathcal{D}'^2}_{\mathcal{D}'^0\mathcal{D}'^1}\ =\ \mathbb{P}\,\Big(\mathop{\oplus}\limits_{(\mathcal{D}^0,\mathcal{D}^1,\mathcal{D}^2)}\mathbb{A}^{\mathcal{D}^1\mathcal{D}^2}_{\mathcal{D}^0\mathcal{D}^1}\Big)\,\widetilde{\mathbb{P}}\Big|_{\mathbb{H}'^{\mathcal{D}'^1}_{\mathcal{D}'^0}\otimes\,\mathbb{H}'^{\mathcal{D}'^2}_{\mathcal{D}'^1}}\tag{7.18}$$

where the operator in parenthesis maps from $\widetilde{\mathbb{H}}'^{\mathcal{D}'^1\mathcal{D}'^2}_{\mathcal{D}'^0\mathcal{D}'^1}$ to $\widetilde{\mathbb{H}}'^{\mathcal{D}'^2}_{\mathcal{D}'^0}$. The projector $\mathbb{P}$ above is spurious, since the result lands anyway in $\mathbb{H}'^{\mathcal{D}'^2}_{\mathcal{D}'^0}$ due to the symmetry property

$$\mathbb{A}^{\mathcal{D}^1\mathcal{D}^2}_{\mathcal{D}^0\mathcal{D}^1}\ \mathbb{U}(z)\otimes\mathbb{U}(z)\ =\ \mathbb{U}(z)\ \mathbb{A}^{\mathcal{D}^1_z\mathcal{D}^2_z}_{\mathcal{D}^0_z\mathcal{D}^1_z}\tag{7.19}$$

of the operator product. Formula (7.18) descends to the multiplicity spaces. Let

$$\widetilde{M}'^{\mathcal{D}'^1\mathcal{D}'^2}_{\mathcal{D}'^0\mathcal{D}'^1\lambda\,\mu}\ =\ \mathop{\oplus}\limits_{(\mathcal{D}^0,\mathcal{D}^1,\mathcal{D}^2)}M^{\mathcal{D}^1}_{\mathcal{D}^0\lambda}\otimes M^{\mathcal{D}^2}_{\mathcal{D}^1\mu}\ \subset\ \widetilde{M}'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda}\otimes\widetilde{M}'^{\mathcal{D}'^2}_{\mathcal{D}'^1\mu}$$

and let $\widetilde{P}$ denote the orthogonal projection from $\widetilde{M}'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda}\otimes\widetilde{M}'^{\mathcal{D}'^2}_{\mathcal{D}'^1\mu}$ to $\widetilde{M}'^{\mathcal{D}'^1\mathcal{D}'^2}_{\mathcal{D}'^0\mathcal{D}'^1\lambda\,\mu}$. Then

$$A'^{\mathcal{D}'^1\mathcal{D}'^2\nu}_{\mathcal{D}'^0\mathcal{D}'^1\lambda\,\mu}\ =\ P_\nu\,\Big(J\otimes\mathop{\oplus}\limits_{\substack{(\lambda^0,\lambda^1,\lambda^2)\\ \lambda^s\in[\lambda^s]}}A^{\lambda^1\lambda^2\nu}_{\lambda^0\lambda^1\lambda\,\mu}\Big)\,\widetilde{P}\Big|_{M'^{\mathcal{D}'^1}_{\mathcal{D}'^0\lambda}\otimes M'^{\mathcal{D}'^2}_{\mathcal{D}'^1\mu}}\,.\tag{7.20}$$

The projector P_ν, see Eq. (7.15), may again be dropped due to the symmetry property (7.19) that on the multiplicity spaces in the realization (7.7) reduces to the relation

$$A^{\lambda^1\lambda^2\nu}_{\lambda^0\lambda^1\lambda\,\mu}\ w_z\otimes w_z\ =\ w_z\otimes I\ A^{\lambda^1_z\lambda^2_z\nu}_{\lambda^0_z\lambda^1_z\lambda\,\mu}\,,\tag{7.21}$$

see (7.17) and (7.10). To further elucidate the above formulae, in Appendix C we work out their details for the case of the $SO(3)$ WZW model.

8 Conclusions

We have discussed in this paper how to set up carefully the Lagrangian approach to the WZW models with boundary using the concepts of gerbes and gerbe modules. The possible boundary conditions involving the Chan-Paton coupling to gauge fields were described in terms of D-branes in the target group carrying gerbe modules. The corresponding gerbes are obtained by the restriction to the D-branes of the gerbe on the target group with the curvature proportional to the invariant 3-form $\mathrm{tr}\,(g^{-1}dg)^3$. In particular, we have discussed the (maximally) symmetric branes that preserve the diagonal current-algebra symmetry of the WZW model with a compact simple group as the target. Such branes are supported by the integrable conjugacy classes in the target group, i.e. the ones that contain elements $\mathrm{e}^{2\pi\mathrm{i}\lambda/\mathsf{k}}$ where k is the level of the model and λ is the (integrable) highest weight of a chiral current algebra primary field.

For a simply connected target group G, up to isomorphism, there is exactly one 1-dimensional gerbe module that leads to a current-algebra symmetric boundary condition over each integrable conjugacy class $\mathcal{C}_\tau\subset G$ with $\mathsf{k}\tau=\lambda$, in agreement with Cardy's classification of boundary conditions [16]. The N-dimensional gerbe modules with the same property are direct sums of the 1-dimensional one. In particular, the symmetric branes in the WZW models with a simply connected target carry only Abelian (twisted) gauge fields.

For the non-simply connected target groups $G'=G/Z$, the integrable conjugacy classes $\mathcal{C}'_{[\tau]}\subset G'$ correspond to Z-orbits $[\tau]$ of rescaled weights τ, with $z\mathcal{C}_\tau\equiv\mathcal{C}_{z\tau}\subset G$ projecting to the same conjugacy class in G' as $\mathcal{C}_\tau\subset G$. The construction of gerbe modules over $\mathcal{C}'_{[\tau]}$ that define symmetric branes required solving a cohomological equation $\delta\mathcal{W}=\mathcal{V}$ with $\mathcal{V}$ an explicit 2-cocycle on Z taking values in the Z-module of $U(1)$-valued functions on the orbit $[\tau]$. The scalar solutions $\mathcal{W}$, if existent, lead to 1-dimensional gerbe modules over $\mathcal{C}'_{[\tau]}$ with non-isomorphic choices labeled by the characters of the fundamental group of $\mathcal{C}'_{[\tau]}$. The N-dimensional gerbe modules are then direct sums of the 1-dimensional ones, resulting again in symmetric branes carrying Abelian gauge fields. The exceptional case involves the groups $Spin(4n)/(\mathbb{Z}_2\times\mathbb{Z}_2)=SO(4n)/\mathbb{Z}_2$ that admit two non-equivalent gerbes differing by discrete torsion. For one of those choices and the integrable conjugacy classes $\mathcal{C}'_{[\tau]}$ corresponding to 1-point orbits $[\tau]$ there are no scalar solutions of the cohomological equation $\delta\mathcal{W}=\mathcal{V}$ and no 1-dimensional gerbe modules leading to current-algebra symmetric boundary conditions. There exist, however, matrix-valued solutions $\mathcal{W}$. The one of minimal rank is provided by the Pauli matrices. It gives rise to a 2-dimensional gerbe module over $\mathcal{C}'_{[\tau]}$. All other gerbe modules resulting in symmetric branes supported by such conjugacy classes are direct sums of the 2-dimensional one. The corresponding branes carry a (projectively flat, twisted) non-Abelian $U(2)$ gauge field. The spontaneous enhancement of the gauge symmetry is even more common in the boundary WZW models with non-simple targets [31, 70]. We plan to return to this subject, and to the geometric classification of the so called twisted-symmetric branes [9] in the WZW models and of branes in general coset theories [38, 26] in the future. Another theme left for future is the geometric treatment along similar lines of the WZW amplitudes on non-oriented worldsheets.

In the last section of the paper we have indicated how the solutions of the cohomological equations defining symmetric branes in the WZW models with non-simply connected targets enter directly the expressions for the boundary partition functions and operator product coefficients of those models. This demonstrated the utility of the geometric considerations which provide a simple and unified view on both classical and quantum WZW theory. The general discussion was illustrated by the detailed computation in Appendix C for the WZW model with the $SO(3)$ group target called, in the algebraic approach, the non-diagonal $D_{\frac{k}{2}+2}$ series of the WZW models with $\widehat{su}_2$ symmetry.

An interesting question left aside concerns the stability [6, 58] of the symmetric branes discussed in this paper, see also Sect. 4.3 of [67]. The natural guess is that, in the present context, stable branes are the ones with gerbe modules of the minimal dimension whose number is equal to the dimension of the bulk Ishibashi states [70]. We plan in the future to address the stability issue and its geometric formulation. This question is also related to that of the Ramond-Ramond charges conserved in the brane condensation processes in the supersymmetric version of the WZW model [72, 10, 28, 11, 12, 35].

Appendix A List of 1-dimensional symmetric $\mathcal{G}'$-branes

==

Covering Group: $\mathbf{SU(r+1)}$

Center: $\cong \mathbb{Z}_{r+1}$

Generator: $z\,, \;\; z^{r+1} = 1$

Action on $\mathcal{A}_W$: $z\tau_i = \tau_{i+1}\,, \;\; i = 0, 1, \ldots, r-1\,, \quad z\tau_r = \tau_0$

weight $\lambda \equiv \mathsf{k}\tau$: $\sum\limits_{i=0}^{r} n_i \tau_i\,, \;\; n_i \in \mathbb{Z}$

Subgroup Z : $\{\, z^n \mid n$ divisible by $N'' = (r+1)/N' \,\} \cong \mathbb{Z}_{N'}$

Quotient group: $SU(r+1)/\mathbb{Z}_{N'}$

---

level: $\mathsf{k} \in \mathbb{Z}$ if N' odd or N'' even,
$\mathsf{k} \in 2\mathbb{Z}$ if N' even and N'' odd

2-**chain** V : $V_{z^n,z^m} \;\equiv\; 1$

2-**cocycle** $\mathcal{V}$:

$$\mathcal{V}_{\tau;z^n,z^m} \;=\; \begin{cases} 1 & \text{if } \; n+m \le r\,, \\ (-1)^{r\sum\limits_{i=0}^{r} i n_i} = \mathrm{e}^{\pi \mathrm{i}\, r(r+1)\, Q_z(\lambda)} & \text{if } \; n+m > r \end{cases}$$

for n, m divisible by N'', $0 \le n, m \le r$

Stabilizers: $Z_{[\tau]} \cong \mathbb{Z}_{n'}$ if n' is the largest integer dividing N' such that $n_i = n_j$ for $j = i + (r+1)/n' \bmod (r+1)$,
possible if n' divides k

Special solution: $\mathcal{W}^{\widehat{\tau}}_{\tau;z^n} \;=\; \mathrm{e}^{\pi \mathrm{i}\, \sigma\, n/(r+1)}$
where $\sigma = 0, 1$ is the parity of $r\sum\limits_{i=0}^{r} i n_i = r(r+1)\, Q_z(\lambda) \bmod 2$
constant on $[\tau]$

cocycle $\widehat{\phi}$: $\widehat{\phi}_{\widehat{\tau};z^n}(z_0) \;\equiv\; 1$

General solution: $\mathcal{W}_{\tau;z^n} \;=\; \mathrm{e}^{2\pi \mathrm{i} n p/(r+1)}\, \mathcal{W}^{\widehat{\tau}}_{\tau;z^n}$ for $p = 0, 1, \ldots, n'-1$
if $Z_{[\tau]} \cong \mathbb{Z}_{n'}$

Covering group: Spin(2r + 1)

Center: $\cong \mathbb{Z}_2$

Generator: $z\,,\ \ z^2 = 1$

Action on $\mathcal{A}_W$: $z\tau_0 = z\tau_1\,, \quad z\tau_1 = \tau_0\,, \quad z\tau_i = \tau_i\,,\ \ i = 2, \ldots, r$

weight $\lambda \equiv \mathsf{k}\tau$: $\sum\limits_{i=0}^{r} n_i\tau_i\,,\ \ n_0, n_1, n_r \in \mathbb{Z},\ \ n_i \in 2\mathbb{Z}$ for $i = 2, \ldots, r-1$

Subgroup Z : $\{1, z\} \cong \mathbb{Z}_2$

Quotient group: $SO(2r+1)$

---

level: $\mathsf{k} \in \mathbb{Z}$

2-**chain V :** $V_{z^n,z^m} \ \equiv\ 1$

2-**cocycle $\mathcal{V}$:** $\mathcal{V}_{\tau;z^n,z^m} \ = \ \begin{cases} 1 & \text{if} \ \ n+m \le 1\,, \\ (-1)^{rn_r} = \mathrm{e}^{2\pi \mathrm{i}\, r\, Q_z(\lambda)} & \text{if} \ \ n+m = 2 \end{cases}$

for $n, m = 0, 1$

Stabilizers: $Z_{[\tau]} = \{1, z\}$ if $n_0 = n_1$ and $Z_{[\tau]} = \{1\}$ otherwise

Special solution: $\mathcal{W}^{\widehat{\tau}}_{\tau;z^n} \ = \ \mathrm{e}^{\pi \mathrm{i} \sigma n/2}$

where $\sigma = 0, 1$ is the parity of $rn_r = 2r\, Q_z(\lambda) \bmod 2$

constant on $[\tau]$

cocycle $\widehat{\phi}$: $\widehat{\phi}_{\widehat{\tau};z^n}(z_0) \ \equiv\ 1$

General solution: $\mathcal{W}_{\tau;z^n} \ = \ (-1)^{np}\, \mathcal{W}^{\widehat{\tau}}_{\tau;z^n}$ for $p = 0, 1$ if $Z_{[\tau]} = \{1, z\}$

Covering group: Sp(2r)

Center: $\cong \mathbb{Z}_2$

Generator: $z\,,\ \ z^2 = 1$

Action on $\mathcal{A}_W$ **:** $z\tau_i = \tau_{r-i}\,,\ \ i = 0, \ldots, r$

weight $\lambda \equiv \mathsf{k}\tau$ **:** $\sum\limits_{i=0}^{r} n_i \tau_i\,,\ \ n_i \in \mathbb{Z}$ for $i = 0, \ldots, r$

Subgroup Z **:** $\{1, z\} \cong \mathbb{Z}_2$

Quotient group: $Sp(2r)/\mathbb{Z}_2$

---

level: $\mathsf{k} \in \mathbb{Z}$ if r even and $\mathsf{k} \in 2\mathbb{Z}$ if r is odd

2-**chain** V **:** $\mathcal{V}_{\tau;z^n,z^m} \ \equiv\ 1$

2-**cocycle** $\mathcal{V}$ **:**

$$\mathcal{V}_{\tau;z^n,z^m} \ = \ \begin{cases} 1 & \text{if} \ \ n+m \le 1\,, \\ (-1)^{\sum\limits_{i=0}^{r} i n_i} = \mathrm{e}^{2\pi \mathrm{i}\, Q_z(\lambda)} & \text{if} \ \ n+m = 2 \end{cases}$$

for $n, m = 0, 1$

Stabilizers: $Z_{[\tau]} = \{1, z\}$ if $n_i = n_{r-i},\ i = 0, \ldots, r,$
$Z_{[\tau]} = \{1\}$ otherwise

Special solution: $\mathcal{W}^{\widehat{\tau}}_{\tau;z^n} \ = \ \mathrm{e}^{\pi \mathrm{i} \sigma n/2}$

where $\sigma = 0, 1$ is the parity of $\sum\limits_{i=0}^{r} i n_i = 2\, Q_z(\lambda) \bmod 2$

constant on $[\tau]$

cocycle $\widehat{\phi}$ **:** $\widehat{\phi}_{\widehat{\tau};z^n}(z_0) \ \equiv\ 1$

General solution: $\mathcal{W}_{\tau;z^n} \ = \ (-1)^{np}\, \mathcal{W}^{\widehat{\tau}}_{\tau;z^n}$ for $p = 0, 1$ if $Z_{[\tau]} = \{1, z\}$

Covering group: Spin(2r) for odd r

Center: $\cong \mathbb{Z}_4$

Generator: $z\,,\;\; z^4 = 1$

Action on $\mathcal{A}_W$: $z\tau_0 = \tau_{r-1}\,, \quad z\tau_1 = \tau_r\,, \quad z\tau_i = \tau_{r-i}\,,\;\; i = 2,\dots,r$

weight $\lambda \equiv \mathsf{k}\tau$ **:** $\sum\limits_{i=0}^{r} n_i\tau_i\,, \;\; n_0, n_1, n_{r-1}, n_r \in \mathbb{Z}\,,$
$n_i \in 2\mathbb{Z}$ for $i = 2,\dots,r-2$

Subgroup Z **:** $\{1, z^2\} \cong \mathbb{Z}_2$

Quotient group: $SO(2r)$

level: $\mathsf{k} \in \mathbb{Z}$

2-**chain** V **:** $V_{z^{2n},z^{2m}} \;\equiv\; 1$

2-**cocycle** $\mathcal{V}$ **:**

$$\mathcal{V}_{\tau;\,z^{2n},z^{2m}} \;=\; \begin{cases} 1 & \text{if} \;\; n+m \le 1\,, \\ (-1)^{n_{r-1}+n_r} = \mathrm{e}^{4\pi\mathrm{i}\, Q_z(\lambda)} & \text{if} \;\; n+m = 2 \end{cases}$$

for $n, m = 0, 1$

Stabilizers:
$Z_{[\tau]} = \{1, z^2\}$ if $n_0 = n_1$ and $n_{r-1} = n_r$, possible if $\mathsf{k} \in 2\mathbb{Z}$,
$Z_{[\tau]} = \{1\}$ otherwise

Special solution: $\mathcal{W}^{\widehat{\tau}}_{\tau;\,z^{2n}} \;=\; \mathrm{e}^{\pi\mathrm{i}\sigma n/2}$
where $\sigma = 0, 1$ is the parity of $n_{r-1} + n_r = 4\,Q_z(\lambda) \bmod 2$
constant on $[\tau]$

cocycle $\widehat{\phi}$**:** $\widehat{\phi}_{\widehat{\tau};\,z^{2n}}(z_0) \;\equiv\; 1$

General solution: $\mathcal{W}_{\tau;\,z^{2n}} \;=\; (-1)^{np}\, \mathcal{W}^{\widehat{\tau}}_{\tau;\,z^{2n}}$ for $p = 0, 1$ if $Z_{[\tau]} = \{1, z^2\}$

– –

Subgroup Z: $\cong \mathbb{Z}_4$

Quotient group: $SO(2r)/\mathbb{Z}_2$

– –

level: $\mathsf{k} \in 2\mathbb{Z}$

2-**chain** V: $V_{z^n,z^m} \;\equiv\; 1$

2-**cocycle** $\mathcal{V}$: $\mathcal{V}_{\tau;z^n,z^m} \;=\; \begin{cases} 1 & \text{if } \; n+m<4\,, \\ (-1)^{n_{r-1}+n_r} = \mathrm{e}^{4\pi\mathrm{i}\,Q_z(\lambda)} & \text{if } \; n+m\geq 4 \end{cases}$

for $n,m=0,1,2,3$

Stabilizers:

$Z_{[\tau]} \cong \mathbb{Z}_4$ — if $n_0=n_1=n_{r-1}=n_r$ and $n_i=n_{r-i},\ i=2,\dots,r-2$, possible if $\mathsf{k}\in 4\mathbb{Z}$,

$Z_{[\tau]} = \{1,z^2\} \cong \mathbb{Z}_2$ — if $n_0=n_1$ and $n_{r-1}=n_r$ but $Z_{[\tau]} \not\cong \mathbb{Z}_4$,

$Z_{[\tau]} = \{1\}$ — otherwise

Special solution: $\mathcal{W}^{\widehat{\tau}}_{\tau;z^n} \;=\; \mathrm{e}^{\pi\mathrm{i}\sigma n/4}$

where $\sigma=0,1$ is the parity of $n_{r-1}+n_r = 4\,Q_z(\lambda) \bmod 2$

constant on $[\tau]$

cocycle $\widehat{\phi}$: $\widehat{\phi}_{\widehat{\tau};z^n}(z_0) \;\equiv\; 1$

General solution: $\mathcal{W}_{\tau;z^n} \;=\; \mathrm{e}^{\pi\mathrm{i}pn/2}\,\mathcal{W}^{\widehat{\tau}}_{\tau;z^n}$ for $p=0,\dots,n'-1$ if $Z_{[\tau]}\cong\mathbb{Z}_{n'}$

Covering group: **Spin(2r)** for even r

Center: $\cong \mathbb{Z}_2 \times \mathbb{Z}_2$

Generators: $z_1,\, z_2\,,\ \ z_1^2 = 1 = z_2^2$

Action on $\mathcal{A}_W$**:** $z_1\tau_i = \tau_{r-i}\,,\ i = 0,\dots,r\,,$

$z_2\tau_0 = \tau_1\,,\quad z_2\tau_1 = \tau_0\,,\quad z_2\tau_i = \tau_i\,,\ i = 2,\dots,r-2\,,$

$z_2\tau_{r-1} = \tau_r\,,\quad z_2\tau_r = \tau_{r-1}$

weight $\lambda \equiv \mathsf{k}\tau$**:** $\sum\limits_{i=0}^{r} n_i\tau_i\,,\ \ n_0, n_1, n_{r-1}, n_r \in \mathbb{Z}\,,$

$n_i \in 2\mathbb{Z}$ for $i = 2,\dots,r-2$

Subgroup Z**:** $\{1, z_1\} \cong \mathbb{Z}_2$

Quotient group: $Spin(2r)/\{1, z_1\}$

level: $\mathsf{k} \in \mathbb{Z}$ if $r/2$ is even,

$\mathsf{k} \in 2\mathbb{Z}$ if $r/2$ is odd

2-**chain** V**:** $V_{z_1^n, z_1^m} \ \equiv\ 1$

2-**cocycle** $\mathcal{V}$**:**

$$\mathcal{V}_{\tau; z_1^n, z_1^m} \ = \begin{cases} 1 & \text{if} \ \ n+m \le 1\,, \\ (-1)^{\frac{r}{2}(n_{r-1}+n_r)} = \mathrm{e}^{\pi \mathrm{i}\, r\, Q_{z_2}(\lambda)} & \text{if} \ \ n+m = 2 \end{cases}$$

for $n, m = 0, 1$

Stabilizers: $Z_{[\tau]} = \{1, z_1\}$ if $n_i = n_{r-i},\ i = 0,\dots,r,$ possible if $\mathsf{k} \in 2\mathbb{Z}$,

$Z_{[\tau]} = \{1\}$ otherwise

Special solution: $\mathcal{W}^{\widehat{\tau}}_{\tau; z_1^n} \ = \ \mathrm{e}^{\pi \mathrm{i} \sigma n/2}$

where $\sigma = 0, 1$ is the parity of $\frac{r}{2}(n_{r-1} + n_r) = r\, Q_{z_2}(\lambda) \bmod 2$ constant on $[\tau]$

cocycle $\widehat{\phi}$**:** $\widehat{\phi}_{\widehat{\tau}; z_1^n}(z_0) \ \equiv\ 1$

General solution: $\mathcal{W}_{\tau; z_1^n} \ = \ (-1)^{np}\, \mathcal{W}^{\widehat{\tau}}_{\tau; z_1^n}$ for $p = 0, 1$ if $Z_{[\tau]} = \{1, z_1\}$

Subgroup Z : $\{1, z_2\} \cong \mathbb{Z}_2$

Quotient group: $SO(2r)$

level: $\mathsf{k} \in \mathbb{Z}$

2-**chain** V : $V_{z_2^n, z_2^m} \;\equiv\; 1$

2-**cocycle** $\mathcal{V}$:

$$\mathcal{V}_{\tau; z_2^n, z_2^m} \;=\; \begin{cases} 1 & \text{if} \;\; n+m \leq 1, \\ (-1)^{n_{r-1}+n_r} = \mathrm{e}^{2\pi \mathrm{i}\, Q_{z_2}(\lambda)} & \text{if} \;\; n+m = 2 \end{cases}$$

for $n, m = 0, 1$

Stabilizers:

$Z_{[\tau]} = \{1, z_2\}$ if $n_0 = n_1$ and $n_{r-1} = n_r$, possible if $\mathsf{k} \in 2\mathbb{Z}$,

$Z_{[\tau]} = \{1\}$ otherwise

Special solution: $\mathcal{W}^{\widehat{\tau}}_{\tau; z_2^n} \;=\; \mathrm{e}^{\pi \mathrm{i} \sigma n/2}$

where $\sigma = 0, 1$ is the parity of $n_{r-1} + n_r = 2\, Q_{z_2}(\lambda) \bmod 2$ constant on $[\tau]$

cocycle $\widehat{\phi}$: $\widehat{\phi}_{\widehat{\tau}; z_2^n}(z_0) \;\equiv\; 1$

General solution: $\mathcal{W}_{\tau; z_2^n} \;=\; (-1)^{np}\, \mathcal{W}^{\widehat{\tau}}_{\tau; z_2^n}$ for $p = 0, 1$ if $Z_{[\tau]} = \{1, z_2\}$

– –

Subgroup Z: $\{1, z_1z_2\} \cong \mathbb{Z}_2$

Quotient group: $Spin(2r)/\{1, z_1z_2\}$

– –

level: $\mathsf{k} \in \mathbb{Z}$ if $r/2$ is even,
$\mathsf{k} \in 2\mathbb{Z}$ if $r/2$ is odd

2-**chain** V: $V_{(z_1z_2)^n,(z_1z_2)^m} \;\equiv\; 1$

2-**cocycle** $\mathcal{V}$:

$$\mathcal{V}_{\tau;(z_1z_2)^n,(z_1z_2)^m} = \begin{cases} 1 & \text{if } n+m \le 1, \\ (-1)^{\frac{r}{2}(n_{r-1}+n_r)} = \mathrm{e}^{\pi \mathsf{i}\, r\, Q_{z_2}(\lambda)} & \text{if } n+m = 2 \end{cases}$$

for $n, m = 0, 1$

Stabilizers: $Z_{[\tau]} = \{1, z_1z_2\}$ if $n_0 = n_{r-1}$, $n_1 = n_r$,
and $n_i = n_{r-i}$ for $i = 2, \dots, r-2$,
possible if $\mathsf{k} \in 2\mathbb{Z}$,
$Z_{[\tau]} = \{1\}$ otherwise

Special solution: $\mathcal{W}^{\widehat{\tau}}_{\tau;(z_1z_2)^n} \;=\; \mathrm{e}^{\pi \mathsf{i} \sigma n/2}$

where $\sigma = 0, 1$ is the parity of $\frac{r}{2}(n_{r-1}+n_r) = r\, Q_{z_2}(\lambda) \bmod 2$
constant on $[\tau]$

cocycle $\widehat{\phi}$: $\widehat{\phi}_{\widehat{\tau};(z_1z_2)^n}(z_0) \;\equiv\; 1$

General solution: $\mathcal{W}_{\tau;(z_1z_2)^n} = (-1)^{np}\, \mathcal{W}^{\widehat{\tau}}_{\tau;(z_1z_2)^n}$ for $p = 0, 1$ if $Z_{[\tau]} = \{1, z_1z_2\}$

Subgroup Z: $\{1, z_1, z_2, z_1z_2\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$

Quotient group: $SO(2r)/\mathbb{Z}_2$

level: $\mathsf{k} \in \mathbb{Z}$ if $r/2$ is even,
$\mathsf{k} \in 2\mathbb{Z}$ if $r/2$ is odd

2-**chain** V:
k odd

$$V_{z,z'} = \begin{cases} \mp\mathsf{i} & \text{if } \ (z,z') = (z_2, z_1)\,, \\ \pm\mathsf{i} & \text{if } \ (z,z') = (z_2, z_1z_2), (z_1z_2, z_1), \\ & \qquad\qquad\;\; (z_1z_2, z_1z_2)\,, \\ -1 & \text{if } \ (z,z') = (z_1z_2, z_2)\,, \\ 1 & \text{otherwise} \end{cases}$$

2-**chain** V:
k even

$$V_{z,z'} = \begin{cases} \pm 1 & \text{if } \ (z,z') = (z_2, z_1), (z_2, z_1z_2), \\ & \qquad\qquad\;\; (z_1z_2, z_1), (z_1z_2, z_1z_2), \\ 1 & \text{otherwise} \end{cases}$$

2-**cocycle** $\mathcal{V}$:
k odd

$$\mathcal{V}_{\tau;z,z'} = \begin{cases} \mp\mathsf{i}(-1)^{n_{r-1}+n_r} & \text{if } \ (z,z') = (z_2, z_1)\,, \\ (-1)^{n_{r-1}+n_r} & \text{if } \ (z,z') = (z_2, z_2)\,, \\ \pm\mathsf{i} & \text{if } \ (z,z') = (z_2, z_1z_2), (z_1z_2, z_1z_2)\,, \\ \pm\mathsf{i}(-1)^{n_{r-1}+n_r} & \text{if } \ (z,z') = (z_1z_2, z_1)\,, \\ -(-1)^{n_{r-1}+n_r} & \text{if } \ (z,z') = (z_1z_2, z_2)\,, \\ 1 & \text{otherwise} \end{cases}$$

2-**cocycle** $\mathcal{V}$:
k even, $r/2$ even

$$\mathcal{V}_{\tau;z,z'} = \begin{cases} \pm(-1)^{n_{r-1}+n_r} & \text{if } \ (z,z') = (z_2, z_1), (z_1z_2, z_1), \\ (-1)^{n_{r-1}+n_r} & \text{if } \ (z,z') = (z_2, z_2), (z_1z_2, z_2), \\ \pm 1 & \text{if } \ (z,z') = (z_2, z_1z_2), (z_1z_2, z_1z_2), \\ 1 & \text{otherwise} \end{cases}$$

2-**cocycle** $\mathcal{V}$:
k even, $r/2$ odd

$$\mathcal{V}_{\tau;z,z'} = \begin{cases} (-1)^{n_{r-1}+n_r} & \text{if } \ (z,z') = (z_1, z_1), (z_1, z_1z_2), \\ & \qquad\qquad\;\; (z_2, z_2), (z_1z_2, z_2), \\ \pm(-1)^{n_{r-1}+n_r} & \text{if } \ (z,z') = (z_2, z_1), (z_1z_2, z_1z_2), \\ \pm 1 & \text{if } \ (z,z') = (z_2, z_1z_2), (z_1z_2, z_1), \\ 1 & \text{otherwise} \end{cases}$$

with $(-1)^{n_{r-1}+n_r} = \mathrm{e}^{2\pi\mathsf{i}\, Q_{z_2}(\lambda)}$ and the signs corresponding to two choices $\mathcal{G}'_\pm$ of the gerbe

---

Stabilizers: $Z_{[\tau]} = \{1\}$ if $n_i \neq n_{r-i}$ for some $i = 0, \ldots, r$
and $n_0 \neq n_1$ or $n_{r-1} \neq n_r$
and $n_0 \neq n_{r-1}$ or $n_1 \neq n_r$ or $n_i \neq n_{r-i}$
for some $i = 2, \ldots, r-2$

Special solution: $\mathcal{W}^{\widehat{\tau}}_{\widehat{\tau_z};z'} \;=\; \mathcal{V}_{\widehat{\tau};z,z'}$

cocycle $\widehat{\phi}$: $\widehat{\phi}_{\widehat{\tau};z}(1) \;\equiv\; 1$

Stabilizers: $Z_{[\tau]} = \{1, z_1\}$ if $n_i = n_{r-i}$ for $i = 0, \ldots, r$ but $n_0 \neq n_1$,
possible if $\mathsf{k} \in 2\mathbb{Z}$

Special solution:
$r/2$ even

$$\mathcal{W}^{\widehat{\tau}}_{\widehat{\tau_z};z'} \;=\; \mathcal{V}_{\widehat{\tau};z,z'}$$

Special solution:
$r/2$ odd

$$\mathcal{W}^{\widehat{\tau}}_{\widehat{\tau_z};z'} \;=\; \begin{cases} \mathrm{e}^{\pi\mathrm{i}\sigma/2} & \text{if} \quad (z,z') = (1,z_1),(1,z_1z_2), \\ & \qquad\qquad (z_1,z_1),(z_1,z_1z_2), \\ \pm\mathrm{e}^{-\pi\mathrm{i}\sigma/2} & \text{if} \quad (z,z') = (z_2,z_1),(z_1z_2,z_1), \\ (-1)^{\sigma} & \text{if} \quad (z,z') = (z_2,z_2),(z_1z_2,z_2), \\ \pm\mathrm{e}^{\pi\mathrm{i}\sigma/2} & \text{if} \quad (z,z') = (z_2,z_1z_2),(z_1z_2,z_1z_2), \\ 1 & \text{otherwise} \end{cases}$$

where $\sigma = 0, 1$ is the parity of $n_{r-1} + n_r = 2\,Q_{z_2}(\lambda) \bmod 2$
constant on $[\tau]$

cocycle $\widehat{\phi}$:

$$\widehat{\phi}_{\widehat{\tau};z}(z_0) \;=\; \begin{cases} \pm(-1)^{\sigma} & \text{if} \;\; (z,z_0) = (z_2,z_1),(z_1z_2,z_1)\,, \\ 1 & \text{otherwise} \end{cases}$$

General solution: $\mathcal{W}_{\tau;z_1^n z_2^m} \;=\; (-1)^{np}\, \mathcal{W}^{\widehat{\tau}}_{\tau;z_1^n z_2^m}$ for $p = 0, 1$

Stabilizers: $Z_{[\tau]} = \{1, z_2\}$ if $n_0 = n_1$ and $n_{r-1} = n_r$
but $n_i \neq n_{r-i}$ for some $i = 0, \ldots, r$,
possible if $\mathsf{k} \in 2\mathbb{Z}$

Special solution:

$$\mathcal{W}^{\widehat{\tau}}_{\widehat{\tau_z};z'} \;=\; \begin{cases} \pm 1 & \text{if} \;\; (z,z') = (1,z_1z_2),(z_1,z_2), \\ & \qquad\qquad (z_2,z_1z_2),(z_1z_2,z_2)\,, \\ 1 & \text{otherwise} \end{cases}$$

cocycle $\widehat{\phi}$:

$$\widehat{\phi}_{\widehat{\tau};z}(z_0) \;=\; \begin{cases} \pm 1 & \text{if} \;\; (z,z_0) = (z_1,z_2),(z_1z_2,z_2)\,, \\ 1 & \text{otherwise} \end{cases}$$

General solution: $\mathcal{W}_{\tau;z_1^n z_2^m} \;=\; (-1)^{mp}\, \mathcal{W}^{\widehat{\tau}}_{\tau;z_1^n z_2^m}$ for $p = 0, 1$

– – – – – – – – – – – – – – – – – – – –

Stabilizers: $Z_{[\tau]} = \{1, z_1 z_2\}$ if $n_0 = n_{r-1} \neq n_1 = n_r$
and $n_i = n_{r-i}$ for $i = 2, \ldots, r-2$,
possible if $\mathsf{k} \in 2\mathbb{Z}$

Special solution:
$r/2$ even

$$\mathcal{W}^{\widehat{\tau}}_{\widehat{\tau}_z; z'} = \begin{cases} \left\{ {1 \atop \mathsf{i}} \right\} & \text{if} \quad (z,z') = (1, z_1z_2), (z_1, z_2), \\ & \qquad\qquad (z_2, z_2), (z_1z_1, z_1z_2), \\ (-1)^\sigma \left\{ {1 \atop -\mathsf{i}} \right\} & \text{if} \quad (z,z') = (1, z_2), (z_1, z_1z_2), \\ & \qquad\qquad (z_2, z_1z_2), (z_1z_2, z_2), \\ 1 & \text{otherwise} \end{cases}$$

Special solution:
$r/2$ odd

$$\mathcal{W}^{\widehat{\tau}}_{\widehat{\tau}_z; z'} = \begin{cases} \mathrm{e}^{\pi \mathsf{i} \sigma/2} \left\{ {1 \atop -\mathsf{i}} \right\} & \text{if} \quad (z,z') = (1, z_2), (z_1z_2, z_2), \\ \mathrm{e}^{\pi \mathsf{i} \sigma/2} \left\{ {1 \atop \mathsf{i}} \right\} & \text{if} \quad (z,z') = (1, z_1z_2), (z_1, z_2), \\ & \qquad\qquad (z_2, z_2), (z_1z_2, z_1z_2), \\ (-1)^\sigma & \text{if} \quad (z,z') = (z_1, z_1), (z_2, z_1), \\ \mathrm{e}^{-\pi \mathsf{i} \sigma/2} \left\{ {1 \atop -\mathsf{i}} \right\} & \text{if} \quad (z,z') = (z_1, z_1z_2), (z_2, z_1z_2), \\ 1 & \text{otherwise} \end{cases}$$

where $\sigma = 0, 1$ is the parity of $n_{r-1} + n_r = 2\, Q_{z_2}(\lambda) \bmod 2$
constant on $[\tau]$

cocycle $\widehat{\phi}$:

$$\widehat{\phi}_{\widehat{\tau}; z}(z_0) \;=\; \begin{cases} \pm(-1)^\sigma & \text{if} \;\; (z, z_0) = (z_1, z_1z_2), (z_2, z_1z_2)\,, \\ 1 & \text{otherwise} \end{cases}$$

General solution:

$$\mathcal{W}_{\tau; z_1^n z_2^m} \;=\; (-1)^{np}\, \mathcal{W}^{\widehat{\tau}}_{\tau; z_1^n z_2^m} \qquad \text{for} \quad p = 0, 1$$

– – – – – – – – – – – – – – – – – – – –

Stabilizers: $Z_{[\tau]} = \{1, z_1, z_2, z_1 z_2\}$ if $n_0 = n_1 = n_{r-1} = n_r$
and $n_i = n_{r-i}$ for $i = 2, \ldots, r-2$,
possible if $\mathsf{k} \in 2\mathbb{Z}$

Special solution:

$$\mathcal{W}^{\widehat{\tau}}_{\tau; z, z'} \;=\; \begin{cases} 1\,, \\ \text{no solution for the lower sign case} \end{cases}$$

cocycle $\widehat{\phi}$:

$$\widehat{\phi}_{\widehat{\tau}; z}(z_0) \;\equiv\; 1$$

General solution:

$$\mathcal{W}_{\tau; z_1^n z_2^m} \;=\; (-1)^{np+mq}\, \mathcal{W}^{\widehat{\tau}}_{\tau; z_1^n z_2^m} \qquad \text{for} \quad p, q = 0, 1$$

Covering group: $\mathbf{E_6}$

Center: $\cong \mathbb{Z}_3$

Generator: $z\,, \;\; z^3 = 1$

Action on $\mathcal{A}_W$: $z\tau_0 = \tau_1\,, \;\; z\tau_1 = \tau_5\,, \;\; z\tau_2 = \tau_4\,, \;\; z\tau_3 = \tau_3\,,$
$z\tau_4 = \tau_6\,, \;\; z\tau_5 = \tau_0\,, \;\; z\tau_6 = \tau_2$

weight $\lambda \equiv \mathsf{k}\tau$: $\sum\limits_{i=0}^{r} n_i \tau_i\,, \;\; n_0, n_1, n_5 \in \mathbb{Z}, \;\; n_2, n_4, n_6 \in 2\mathbb{Z}, \;\; n_3 \in 3\mathbb{Z}$

Subgroup Z : $\{1, z, z^2\} \cong \mathbb{Z}_3$

Quotient group: $E_6/\mathbb{Z}_3$

level: $\mathsf{k} \in \mathbb{Z}$

2-**chain** V : $V_{z^n,z^m} \;\equiv\; 1$

2-**cocycle** $\mathcal{V}$: $\mathcal{V}_{\tau;z^n,z^m} \;=\; 1$

Stabilizers: $Z_{[\tau]} = \{1, z, z^2\}$ if $n_0 = n_1 = n_5$ and $n_2 = n_4 = n_6$,
possible if $\mathsf{k} \in 3\mathbb{Z}$
$Z_{[\tau]} = \{1\}$ otherwise

Special solution: $\mathcal{W}^{\widehat{\tau}}_{\tau;z^n} \;=\; 1$

cocycle $\widehat{\phi}$: $\widehat{\phi}_{\widehat{\tau};z}(z_0) \;\equiv\; 1$

General solution: $\mathcal{W}_{\tau;z^n} \;=\; \mathrm{e}^{2\pi\mathrm{i}np/3}$ for $p = 0, 1, 2$ if $Z_{[\tau]} = \{1, z, z^2\}$

Covering group: $\mathbf{E_7}$

Center: $\cong \mathbb{Z}_2$

Generator: $z\,,\;\; z^2 = 1$

Action on $\mathcal{A}_W$: $z\tau_0 = \tau_1\,,\;\; z\tau_1 = \tau_0\,,\;\; z\tau_i = \tau_{8-i}$ for $i = 2,\ldots,6,\;\; z\tau_7 = \tau_7$

weight $\lambda \equiv \mathsf{k}\tau$: $\sum\limits_{i=0}^{r} n_i\tau_i\,,\;\; n_0, n_1 \in \mathbb{Z},\;\; n_2, n_6, n_7 \in 2\mathbb{Z},$
$n_3, n_5 \in 3\mathbb{Z},\;\; n_4 \in 4\mathbb{Z}$

Subgroup Z: $\{1, z\} \cong \mathbb{Z}_2$

Quotient group: $E_7/\mathbb{Z}_2$

level: $\mathsf{k} \in 2\mathbb{Z}$

2-**chain** V: $V_{z^n,z^m} \;\equiv\; 1$

2-**cocycle** $\mathcal{V}$: $\mathcal{V}_{\tau;z^n,z^m} \;=\; \begin{cases} 1 & \text{if } \; n+m \le 1\,, \\ (-1)^{n_1+n_3+\frac{n_7}{2}} = \mathrm{e}^{2\pi\mathrm{i}\, Q_z(\lambda)} & \text{if } \; n+m = 2 \end{cases}$
for $n, m = 0, 1$

Stabilizers: $Z_{[\tau]} = \{1, z\}$ if $n_0 = n_1,\; n_2 = n_6$ and $n_3 = n_5\,,$
$Z_{[\tau]} = \{1\}$ otherwise

Special solution: $\mathcal{W}^{\widehat{\tau}}_{\tau;z^n} \;=\; \mathrm{e}^{\pi\mathrm{i}\sigma n/2}$

where $\sigma = 0, 1$ is the parity of $n_1 + n_3 + \frac{n_7}{2} = 2\,Q_z(\lambda) \bmod 2$
constant on $[\tau]$

cocycle $\widehat{\phi}$: $\widehat{\phi}_{\widehat{\tau};z^n}(z_0) \;\equiv\; 1$

General solution: $\mathcal{W}_{\tau;z^n} \;=\; (-1)^{np}\, \mathcal{W}^{\widehat{\tau}}_{\tau;z^n}$ for $p = 0, 1$ if $Z_{[\tau]} = \{1, z\}$

Appendix B

We provide here a direct check that the action of the group elements w_z in V_λ intertwines the multiplicity subspaces $M^{\lambda^1}_{\lambda^0\lambda}$, see (7.10). We have to show that under the action of w_z in V_λ an element $|\psi\rangle \in M^{\lambda^1_z}_{\lambda^0_z\lambda} \subset V_\lambda$ lands in $M^{\lambda^1}_{\lambda^0\lambda}$. One has to check then that $w_z|\psi\rangle$ satisfies conditions (7.4), (7.5) and (7.6). The first one follows since

$$\begin{aligned} t\, w_z|\psi\rangle \;=\; w_z\, w_z^{-1} t\, w_z|\psi\rangle \;&=\; [\mathrm{tr}\,(w_z^{-1} t\, w_z)(\lambda^1_z - \lambda^0_z)]\; w_z|\psi\rangle \\ &=\; [\mathrm{tr}\, t(\lambda^1 - \lambda^0)]\; w_z|\psi\rangle\,. \end{aligned}$$

For the second and the third condition, one has to use the relations

$$\begin{aligned} w_z^{-1}\alpha_i\, w_z \quad &= \begin{cases} \alpha_{i_z} & \text{if} \quad i_z \neq 0\,, \\ -\phi & \text{if} \quad i_z = 0\,, \end{cases} \\ w_z^{-1}\phi\, w_z \quad &= -\alpha_{0_z} \end{aligned}$$

easy to check by computing the Killing form between both sides and $\tau_i = \frac{1}{k_i^\vee}\lambda_i$. The above equalities imply that

$$\begin{aligned} w_z^{-1} e_{-\alpha_i}\, w_z \quad &= \begin{cases} e_{-\alpha_{i_z}} & \text{if} \quad i_z \neq 0\,, \\ e_\phi & \text{if} \quad i_z = 0\,, \end{cases} \\ w_z^{-1} e_\phi\, w_z \quad &= e_{-\alpha_{0_z}} \end{aligned}$$

and that

$$\begin{aligned} \mathrm{tr}\,\alpha_i^\vee \lambda^1 \quad &= \begin{cases} \mathrm{tr}\,\alpha_{i_z}\lambda^1_z & \text{if} \quad i_z \neq 0\,, \\ k - \mathrm{tr}\,\phi^\vee\lambda^1_z & \text{if} \quad i_z = 0\,, \end{cases} \\ k - \mathrm{tr}\,\phi^\vee\lambda^1 \quad &= \mathrm{tr}\,\alpha^\vee_{0_z}\lambda^1_z\,. \end{aligned}$$

As a result,

$$\begin{aligned} e_{-\alpha_i}^{\mathrm{tr}\,\alpha_i^\vee\lambda^1+1} w_z|\psi\rangle \quad &= \begin{cases} w_z\, e_{-\alpha_{i_z}}^{\mathrm{tr}\,\alpha^\vee_{i_z}\lambda^1_z+1}|\psi\rangle & \text{if} \quad i_z \neq 0\,, \\ w_z\, e_\phi^{k-\mathrm{tr}\,\phi^\vee\lambda^1_z+1}|\psi\rangle & \text{if} \quad i_z = 0\,, \end{cases} \\ e_\phi^{k-\mathrm{tr}\,\phi^\vee\lambda^1_z+1} w_z|\psi\rangle \quad &= w_z\, e_{-\alpha_{0_z}}^{\mathrm{tr}\,\alpha^\vee_{0_z}\lambda^1_z+1}|\psi\rangle \end{aligned}$$

and the right hand sides vanish since $|\psi\rangle \in M^{\lambda^1_z}_{\lambda^0_z\lambda}$. Note that one of the consequences of the intertwining property of w_z is the symmetry $N^{\lambda^1_z}_{\lambda^0_z\lambda} = N^{\lambda^1}_{\lambda^0\lambda}$ of the fusion coefficients giving the dimensions of the spaces $M^{\lambda^1}_{\lambda^0\lambda}$.

Appendix C

For the sake of illustration, we make explicit the formulae of Sect. 7 for the case of the WZW model with the $SO(3)$ target. With the Cartan subalgebra of $su(2)$ spanned by the Pauli matrix σ^3, we label the weights $j\sigma^3$ integrable at level k by spins $j = 0, \frac{1}{2}, \dots, \frac{\mathsf{k}}{2}$. They correspond to the integrable conjugacy classes $\mathcal{C}_{j\sigma^3/\mathsf{k}} \subset SU(2)$ supporting the rank 1 branes $\mathcal{D}_1$ of the theory with the $SU(2)$ target that we shall also label by j. In the rank 1 case, the multiplicity spaces of the decomposition (7.2) and (7.3) take in the realization (7.7) the form

$$M^{j^1}_{j^0 j} \;=\; \begin{cases} \mathbb{C}\,|j, j^1 - j^0\rangle & \text{if } \; j + j^0 + j^1 \in \mathbb{Z} \;\text{ and }\; |j^0 - j^1| \le j \le \min\big({j^0+j^1 \atop k-j^0-j^1}\big)\,, \\ \{0\} & \text{otherwise} \end{cases}$$

where $|j, m\rangle$ are the vectors of the standard orthonormal bases of the spin j representation V_j labeled by the magnetic number $m = -j, -j+1, \dots, j$. The dimensions $N^{j^1}_{j^0 j}$ are equal 1 or 0. In other words, the multiplicity space $M^{j^1}_{j^0 j}$ is spanned by the vector

$$e^{j^1}_{j^0 j} \;\equiv\; N^{j^1}_{j^0 j}\, |j, j^1 - j^0\rangle \;\in\; V_j\,.$$

The group $SO(3)$ is the quotient of $SU(2)$ by the center $Z = \{1, -1\}$. The level k of the $SO(3)$ model has to be even. The nontrivial element of Z acts on the weights by $j \mapsto \frac{\mathsf{k}}{2} - j \equiv j_-$. One has $N^{j^1_-}_{j^0_- j} = N^{j^1}_{j^0 j}$. We shall label the integrable conjugacy classes in $SO(3)$ by the corresponding Z-orbits of spins, with the two-point orbits $[j] = \{j, j_-\}$, $j = 0, \frac{1}{2}, \dots, \frac{\mathsf{k}}{4} - \frac{1}{2}$, and the single-point one $[\frac{\mathsf{k}}{4}] = \{\frac{\mathsf{k}}{4}\}$. With the choice $w_1 = 1$, $w_{-1} = \big(\begin{smallmatrix} 0 & i \\ i & 0 \end{smallmatrix}\big)$ and $b_{-1,-1} = \frac{1}{2}\sigma^3$ as in [40], the special solution $\mathcal{W}$ of the cohomological equation (5.18), relabeled by spins j such that $\mathsf{k}\tau = j\sigma^3$, has the form

$$\mathcal{W}_{j;1} \;=\; 1\,, \qquad \mathcal{W}_{j;-1} \;=\; \begin{cases} 1 & \text{if } \; 2j \; \text{ is even}\,, \\ \mathsf{i} & \text{if } \; 2j \; \text{ is odd}\,, \end{cases} \tag{C.1}$$

see Appendix A. It induces unique (up to isomorphism) rank 1 branes $\mathcal{D}'_1$ of the $SO(3)$ model supported by the conjugacy classes corresponding to the two-point orbits $[j]$. To simplify notations, we shall label such branes by $[j]$. On the other hand, there are two non-isomorphic choices, induced by $\mathcal{W}_{j;-1}$ of Eq. (C.1) and by its negative, for the branes supported by the conjugacy class corresponding to the single-point orbit $[\frac{\mathsf{k}}{4}]$. We shall label them $[\frac{\mathsf{k}}{4}]_\pm$. Since $w_{-1}\,|j, m\rangle = \mathsf{i}^{2j}\,|j, -m\rangle$, for the solution (C.1) the map

$$U^{j^1}_{j^0 j}(-1) \;:\; M^{j^1_-}_{j^0_- j} \;\longrightarrow\; M^{j^1}_{j^0 j}$$

of Eq. (7.9) takes the form

$$U^{j^1}_{j^0 j}(-1)\; e^{j^1_-}_{j^0_- j} \;=\; u^{j^1}_{j^0 j}\; e^{j^1}_{j^0 j}$$

where

$$u^{j^1}_{j^0 j} \;=\; \begin{cases} \mathsf{i}^{2j} & \text{if } \;\; 2j \;\; \text{ is even}\,, \\ \mathsf{i}^{2j-1} & \text{if } \;\; 2j^0 \;\; \text{ is even and } \;\; 2j^1 \;\; \text{ is odd}\,, \\ \mathsf{i}^{2j+1} & \text{if } \;\; 2j^0 \;\; \text{ is odd and } \;\; 2j^1 \;\; \text{ is even}\,. \end{cases} \tag{C.2}$$

Note that $u^{j^1}_{j^0 j} = u^{j^1}_{j^0_- j} = u^{j^1_-}_{j^0 j}$ and that $(u^{j^1}_{j^0 j})^2 = 1$.

Given operators $U^{j^1}_{j^0 j}(z)$, the multiplicity spaces in the boundary $SO(3)$ model are obtained using relations (7.13), (7.14) and (7.15). First, for $j^0, j^1 < \frac{\mathsf{k}}{4}$, the space $\widetilde{M}'^{[j^1]}_{[j^0]\,j}$ is spanned by the vectors $e^{j^1}_{j^0 j}$, $e^{j^1}_{j^0_- j}$, $e^{j^1_-}_{j^0 j}$, $e^{j^1_-}_{j^0_-}$ so that, with N denoting the diagonal matrix with the entries $N^{j^1}_{j^0 j}$, $N^{j^1}_{j^0_- j}$, $N^{j^1}_{j^0_- j}$, $N^{j^1}_{j^0 j}$, we may identify

$$\widetilde{M}'^{[j^1]}_{[j^0]\,j} \;\cong\; N\,\mathbb{C}^4\,.$$

The operator $U^{[j^1]}_{[j^0]\,j}(-1)$ becomes in this representation a 4×4 anti-diagonal matrix with the entries given by Eq. (C.2). The multiplicity spaces $M'^{[j^1]}_{[j^0]\,j}$ are spanned by the vectors

$$e'^{[j^1]}_{[j^0]\,j} \;=\; \tfrac{1}{\sqrt{2}}\left(e^{j^1}_{j^0 j} + u^{j^1}_{j^0 j}\, e^{j^1_-}_{j^0_- j}\right), \qquad f'^{[j^1]}_{[j^0]\,j} \;=\; \tfrac{1}{\sqrt{2}}\left(e^{j^1}_{j^0_- j} + u^{j^1}_{j^0 j}\, e^{j^1_-}_{j^0 j}\right)$$

and their dimension is

$$N'^{[j^1]}_{[j^0]\,j} \;=\; N^{j^1}_{j^0 j} + N^{j^1}_{j^0_- j}\,.$$

Next, for $j^0 = \frac{\mathsf{k}}{4}$ and $j^1 < \frac{\mathsf{k}}{4}$, the space $\widetilde{M}'^{[j^1]}_{[\frac{\mathsf{k}}{4}]_\pm j}$ is spanned by the vectors $e^{j^1}_{\frac{\mathsf{k}}{4} j}$, $e^{j^1_-}_{\frac{\mathsf{k}}{4} j}$ so that we may identify

$$\widetilde{M}'^{[j^1]}_{[\frac{\mathsf{k}}{4}]_\pm j} \;\cong\; N^{j^1}_{\frac{\mathsf{k}}{4} j}\,\mathbb{C}^2\,.$$

The operator $U^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^0]\,j}(-1)$ is represented by an anti-diagonal 2×2 matrix with the entries given by Eq. (C.2) multiplied by ± 1 for the $[\frac{\mathsf{k}}{4}]_\pm$ branes. The multiplicity space $M'^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^0]\,j}$ is spanned by the vector

$$e'^{[j^1]}_{[\frac{\mathsf{k}}{4}]_\pm j} \;=\; \tfrac{1}{\sqrt{2}}\left(e^{j^1}_{\frac{\mathsf{k}}{4} j} \pm u^{j^1}_{\frac{\mathsf{k}}{4} j}\, e^{j^1_-}_{\frac{\mathsf{k}}{4} j}\right)$$

and has the dimension

$$N'^{[j^1]}_{[\frac{\mathsf{k}}{4}]_\pm j} \;=\; N^{j^1}_{\frac{\mathsf{k}}{4} j}\,.$$

The case $j_0 < \frac{\mathsf{k}}{4}$, $j^1 = \frac{\mathsf{k}}{4}$ is similar with

$$\widetilde{M}'^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^0]\,j} \;\cong\; N^{\frac{\mathsf{k}}{4}}_{j^0 j}\,\mathbb{C}^2\,,$$

the multiplicity space $M'^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^0]\,j}$ spanned by the vector

$$e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^0]\,j} \;=\; \tfrac{1}{\sqrt{2}}\left(e^{\frac{\mathsf{k}}{4}}_{j^0 j} \pm u^{\frac{\mathsf{k}}{4}}_{j^0 j}\, e^{\frac{\mathsf{k}}{4}}_{j^0_- j}\right)$$

and its dimension equal to

$$N'^{\,[\frac{\mathsf{k}}{4}]_\pm}_{[j^0]\,j} \;=\; N^{\frac{\mathsf{k}}{4}}_{j^0 j}\,.$$

Finally, for $j^0 = j^1 = \frac{\mathsf{k}}{4}$, the space $\widetilde{M}'^{\,[\frac{\mathsf{k}}{4}]_\pm}_{[\frac{\mathsf{k}}{4}]_\pm j}$ is spanned by the vector $e^{\frac{\mathsf{k}}{4}}_{\frac{\mathsf{k}}{4} j}$. Note that $N^{\frac{\mathsf{k}}{4}}_{\frac{\mathsf{k}}{4} j}$ is equal to 1 for integer j and to 0 for half-integer j so that $e^{\frac{\mathsf{k}}{4}}_{\frac{\mathsf{k}}{4} j} = 0$ for half-integer j. The operator $U^{[\frac{\mathsf{k}}{4}]_\pm}_{[\frac{\mathsf{k}}{4}]_\pm j}$ acts as multiplication by $(\pm)(\pm)(-1)^j$ with the signs labeling the choices of the branes. The multiplicity space $M'^{\,[\frac{\mathsf{k}}{4}]_\pm}_{[\frac{\mathsf{k}}{4}]_\pm j}$ is spanned by the vector

$$e'^{\,[\frac{\mathsf{k}}{4}]_\pm}_{[\frac{\mathsf{k}}{4}]_\pm j} \;=\; \tfrac{1}{2}\Big(1 + (\pm)(\pm)(-1)^j\Big)\, e^{\frac{\mathsf{k}}{4}}_{\frac{\mathsf{k}}{4} j}$$

and its dimension is

$$N'^{\,[\frac{\mathsf{k}}{4}]_\pm}_{[\frac{\mathsf{k}}{4}]_\pm j} \;=\; \tfrac{1}{2}\Big(1 + (\pm)(\pm)(-1)^j\Big)\, N^{\frac{\mathsf{k}}{4}}_{\frac{\mathsf{k}}{4} j}\,.$$

In particular, for $j = \frac{1}{2}$ and $j^0, j^1 = 0, \frac{1}{2}, \dots, \frac{\mathsf{k}}{4} - \frac{1}{2}$, one obtains:

$$N'^{\,[j^1]}_{[j^0]\frac{1}{2}} \;=\; \begin{cases} 1 & \text{if} \quad |j^0 - j^1| = \frac{1}{2}\,, \\ 0 & \text{otherwise}\,, \end{cases} \qquad N'^{\,[\frac{\mathsf{k}}{4}]_\pm}_{[j^0]\frac{1}{2}} \;=\; \begin{cases} 1 & \text{if} \quad j^0 = \frac{\mathsf{k}}{4} - \frac{1}{2}\,, \\ 0 & \text{otherwise}\,, \end{cases}$$

$$N'^{\,[j^1]}_{[\frac{\mathsf{k}}{4}]_\pm \frac{1}{2}} \;=\; \begin{cases} 1 & \text{if} \quad j^1 = \frac{\mathsf{k}}{4} - \frac{1}{2}\,, \\ 0 & \text{otherwise}\,, \end{cases} \qquad N'^{\,[\frac{\mathsf{k}}{4}]_\pm}_{[\frac{\mathsf{k}}{4}]_\pm \frac{1}{2}} \;=\; 0$$

so that the matrix $N'_{\frac{1}{2}}$ may be identified with the adjacency matrix of the $D_{\frac{\mathsf{k}}{2}+2}$ Dynkin diagram [7].

For the $SU(2)$ theory, the operator product coefficients are given by the map $A^{j^1 j^2 j''}_{j^0 j^1 j\, j'}$ of (7.17) such that

$$A^{j^1 j^2 j''}_{j^0 j^1 j\, j'}\; e^{j^1}_{j^0 j} \otimes e^{j^2}_{j^1 j'} \;=\; F_{j^1 j''}\!\begin{bmatrix} j^0 & j^2 \\ j & j' \end{bmatrix}\; e^{j^2}_{j^0 j''} \otimes e^{j''}_{j\, j'}$$

where the entries of the fusing F-matrix $F_{j^1 j''}\!\begin{bmatrix} j^0 & j^2 \\ j & j' \end{bmatrix}$ vanish unless $N^{j^1}_{j^0 j} N^{j^2}_{j^1 j'} N^{j^2}_{j^0 j''} N^{j''}_{j j'} = 1$. The symmetry (7.21) reduces to the relation

$$F_{j^1_- j''}\!\begin{bmatrix} j^0_- & j^2_- \\ j & j' \end{bmatrix} \;=\; (-1)^{j''-j-j'}\, F_{j^1 j''}\!\begin{bmatrix} j^0 & j^2 \\ j & j' \end{bmatrix}$$

for the fusing matrices, see [13]. For the boundary operator product of the $SO(3)$ model, see [64] for a related work for the minimal models and [52] for the case of level k divisible by 4, one obtains using the relation (7.20) and the identity

$$(-1)^{j''-j-j'}\, u^{j^1}_{j^0 j}\, u^{j^2}_{j^1 j'} \;=\; u^{j^2}_{j^0 j''}$$

holding whenever $N^{j^1}_{j^0 j} N^{j^2}_{j^1 j'} N^{j^2}_{j^0 j''} N^{j''}_{j\,j'} = 1$:

$$
\begin{aligned}
A'^{[j^1][j^2]\,j''}_{[j^0][j^1]\,j\,j'}\;\; e'^{[j^1]}_{[j^0]\,j} \otimes e'^{[j^2]}_{[j^1]\,j'} \quad &= \quad \tfrac{1}{\sqrt{2}}\, F_{j^1 j''}\begin{bmatrix} j^0 & j^2 \\ j & j' \end{bmatrix}\; e'^{[j^2]}_{[j^0]\,j''} \otimes e^{j''}_{j\,j'}\,, \\
A'^{[j^1][j^2]\,j''}_{[j^0][j^1]\,j\,j'}\;\; f'^{[j^1]}_{[j^0]\,j} \otimes f'^{[j^2]}_{[j^1]\,j'} \quad &= \quad \tfrac{1}{\sqrt{2}}\, u^{j^1}_{j^0 j}\; F_{j^1_-\, j''}\begin{bmatrix} j^0 & j^2 \\ j & j' \end{bmatrix}\; e'^{[j^2]}_{[j^0]\,j''} \otimes e^{j''}_{j\,j'}\,, \\
A'^{[j^1][j^2]\,j''}_{[j^0][j^1]\,j\,j'}\;\; e'^{[j^1]}_{[j^0]\,j} \otimes f'^{[j^2]}_{[j^1]\,j'} \quad &= \quad \tfrac{1}{\sqrt{2}}\, u^{j^1}_{j^0 j}\; F_{j^1_-\, j''}\begin{bmatrix} j^0_- & j^2 \\ j & j' \end{bmatrix}\; f'^{[j^2]}_{[j^0]\,j''} \otimes e^{j''}_{j\,j'}\,, \\
A'^{[j^1][j^2]\,j''}_{[j^0][j^1]\,j\,j'}\;\; f'^{[j^1]}_{[j^0]\,j} \otimes e'^{[j^2]}_{[j^1]\,j'} \quad &= \quad \tfrac{1}{\sqrt{2}}\, F_{j^1 j''}\begin{bmatrix} j^0_- & j^2 \\ j & j' \end{bmatrix}\; f'^{[j^2]}_{[j^0]\,j''} \otimes e^{j''}_{j\,j'}\,, \\
A'^{[j^1][j^2]\,j''}_{[\frac{\mathsf{k}}{4}]_\pm[j^1]\,j\,j'}\;\; e'^{[j^1]}_{[\frac{\mathsf{k}}{4}]_\pm\,j} \otimes e'^{[j^2]}_{[j^1]\,j'} \quad &= \quad \tfrac{1}{\sqrt{2}}\, F_{j^1 j''}\begin{bmatrix} \frac{\mathsf{k}}{4} & j^2 \\ j & j' \end{bmatrix}\; e'^{[j^2]}_{[\frac{\mathsf{k}}{4}]_\pm\,j''} \otimes e^{j''}_{j\,j'}\,, \\
A'^{[j^1][j^2]\,j''}_{[\frac{\mathsf{k}}{4}]_\pm[j^1]\,j\,j'}\;\; e'^{[j^1]}_{[\frac{\mathsf{k}}{4}]_\pm\,j} \otimes f'^{[j^2]}_{[j^1]\,j'} \quad &= \quad \pm\tfrac{1}{\sqrt{2}}\, u^{j^1}_{\frac{\mathsf{k}}{4}\,j}\; F_{j^1_-\, j''}\begin{bmatrix} \frac{\mathsf{k}}{4} & j^2 \\ j & j' \end{bmatrix}\; e'^{[j^2]}_{[\frac{\mathsf{k}}{4}]_\pm\,j''} \otimes e^{j''}_{j\,j'}\,, \\
A'^{[\frac{\mathsf{k}}{4}]_\pm[j^2]\,j''}_{[j^0][\frac{\mathsf{k}}{4}]_\pm\,j\,j'}\;\; e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^0]\,j} \otimes e'^{[j^2]}_{[\frac{\mathsf{k}}{4}]_\pm\,j'} \quad &= \quad \tfrac{1}{\sqrt{2}}\, F_{\frac{\mathsf{k}}{4}\,j''}\begin{bmatrix} j^0 & j^2 \\ j & j' \end{bmatrix}\; e'^{[j^2]}_{[j^0]\,j''} \otimes e^{j''}_{j\,j'} \\
&\quad\;\; \pm\, \tfrac{1}{\sqrt{2}}\, u^{\frac{\mathsf{k}}{4}}_{j^0 j}\; F_{\frac{\mathsf{k}}{4}\,j''}\begin{bmatrix} j^0_- & j^2 \\ j & j' \end{bmatrix}\; f'^{[j^2]}_{[j^0]\,j''} \otimes e^{j''}_{j\,j'}\,, \\
A'^{[j^1][\frac{\mathsf{k}}{4}]_\pm\,j''}_{[j^0][j^1]\,j\,j'}\;\; e'^{[j^1]}_{[j^0]\,j} \otimes e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^1]\,j'} \quad &= \quad \tfrac{1}{\sqrt{2}}\, F_{j^1 j''}\begin{bmatrix} j^0 & \frac{\mathsf{k}}{4} \\ j & j' \end{bmatrix}\; e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^0]\,j''} \otimes e^{j''}_{j\,j'}\,, \\
A'^{[j^1][\frac{\mathsf{k}}{4}]_\pm\,j''}_{[j^0][j^1]\,j\,j'}\;\; f'^{[j^1]}_{[j^0]\,j} \otimes e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^1]\,j'} \quad &= \quad \pm\, \tfrac{1}{\sqrt{2}}\, u^{\frac{\mathsf{k}}{4}}_{j^0 j''}\; F_{j^1 j''}\begin{bmatrix} j^0_- & \frac{\mathsf{k}}{4} \\ j & j' \end{bmatrix}\; e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^0]\,j''} \otimes e^{j''}_{j\,j'}\,, \\
A'^{[\frac{\mathsf{k}}{4}]_\pm[j^2]\,j''}_{[\frac{\mathsf{k}}{4}]_\pm[\frac{\mathsf{k}}{4}]_\pm\,j\,j'}\;\; e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[\frac{\mathsf{k}}{4}]_\pm\,j} \otimes e'^{[j^2]}_{[\frac{\mathsf{k}}{4}]_\pm\,j'} \quad &= \quad \tfrac{1}{2}\Big(1 + (\pm)(\pm)(-1)^{j}\Big) \\
&\quad\;\; \cdot\, F_{\frac{\mathsf{k}}{4}\,j''}\begin{bmatrix} \frac{\mathsf{k}}{4} & j^2 \\ j & j' \end{bmatrix}\; e'^{[j^2]}_{[\frac{\mathsf{k}}{4}]_\pm\,j''} \otimes e^{j''}_{j\,j'}\,, \\
A'^{[\frac{\mathsf{k}}{4}]_\pm[\frac{\mathsf{k}}{4}]_\pm\,j''}_{[j^0][\frac{\mathsf{k}}{4}]_\pm\,j\,j'}\;\; e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^0]\,j} \otimes e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[\frac{\mathsf{k}}{4}]_\pm\,j'} \quad &= \quad \tfrac{1}{2}\Big(1 + (\pm)(\pm)(-1)^{j'}\Big) \\
&\quad\;\; \cdot\, F_{\frac{\mathsf{k}}{4}\,j''}\begin{bmatrix} j^0 & \frac{\mathsf{k}}{4} \\ j & j' \end{bmatrix}\; e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^0]\,j''} \otimes e^{j''}_{j\,j'}\,, \\
A'^{[j^1][\frac{\mathsf{k}}{4}]_\pm\,j''}_{[\frac{\mathsf{k}}{4}]_\pm[j^1]\,j\,j'}\;\; e'^{[j^1]}_{[\frac{\mathsf{k}}{4}]_\pm\,j} \otimes e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[j^1]\,j'} \quad &= \quad F_{j^1 j''}\begin{bmatrix} \frac{\mathsf{k}}{4} & \frac{\mathsf{k}}{4} \\ j & j' \end{bmatrix}\; e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[\frac{\mathsf{k}}{4}]_\pm\,j''} \otimes e^{j''}_{j\,j'}\,, \\
A'^{[\frac{\mathsf{k}}{4}]_\pm[\frac{\mathsf{k}}{4}]_\pm\,j''}_{[\frac{\mathsf{k}}{4}]_\pm[\frac{\mathsf{k}}{4}]_\pm\,j\,j'}\;\; e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[\frac{\mathsf{k}}{4}]_\pm\,j} \otimes e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[\frac{\mathsf{k}}{4}]_\pm\,j'} \quad &= \quad \tfrac{1}{2}\Big(1 + (\pm)(\pm)(-1)^{j}\Big) \\
&\quad\;\; \cdot\, F_{\frac{\mathsf{k}}{4}\,j''}\begin{bmatrix} \frac{\mathsf{k}}{4} & \frac{\mathsf{k}}{4} \\ j & j' \end{bmatrix}\; e'^{[\frac{\mathsf{k}}{4}]_\pm}_{[\frac{\mathsf{k}}{4}]_\pm\,j''} \otimes e^{j''}_{j\,j'}\,.
\end{aligned}
$$

References

[1] Alekseev, A., Meinrenken, E., Woodward, C.: *The Verlinde formulas as fixed point formulas.* J. Symplectic Geom. **1** (2001), 1-46 and **1** (2002), 427-434

[2] Alekseev, A. Yu., Schomerus, V.: *D-branes in the WZW model.* Phys. Rev. **D 60** (1999), R061901-R061902

[3] Alvarez, O.: *Topological quantization and cohomology.* Commun. Math. Phys. **100** (1985), 279-309

[4] Alvarez-Gaumé, L., Gomez, C., Sierra, G.: *Quantum group interpretation of some conformal field theories.* Phys. Lett. **B 220** (1989), 142-152

[5] Aspinwall, P., S.: *D-branes on Calabi-Yau manifolds.* arXiv:hep-th/0403166

[6] Bachas, C., Douglas, M. Schweigert, C.: *Flux stabilization of D-branes.* J. High Energy Phys. **JHEP05** (2000), 048

[7] Behrend, R. E., Pearce, P. A., Petkova, V. B., Zuber, J.-B.: *Boundary conditions in rational conformal field theories.* Nucl. Phys. **B 579** (2000), 707-773

[8] Behrend, K., Xu, P. Zhang, B.: *Equivariant gerbes over compact simple Lie groups.* C. R. Acad. Sci. Paris **336** Sér. I (2003), 251-256

[9] Birke, L., Fuchs, J., Schweigert, C.: *Symmetry breaking boundary conditions and WZW orbifolds.* Adv. Theor. Math. Phys. **3** (1999), 671-726

[10] Bouwknegt, P., Carey, A. L., Mathai, V., Murray, M. K., Stevenson, D., *Twisted K-theory and K-theory of bundle gerbes.* Comm. Math. Phys. **228** (2002), 17-45

[11] Bouwknegt, P., Dawson, P., Ridout, D.: *D-branes on group manifolds and fusion rings.* J. High Energy Phys. **JHEP12** (2002), 065

[12] Bouwknegt, P., Ridout, D.: *A note on the equality of algebraic and geometric D-brane charges in WZW models.* J. High Energy Phys. **JHEP05** (2004), 029

[13] Bruner, I., Schomerus, V.: *On Superpotentials for D-branes in Gepner models.* J. High Energy Phys. **JHEP10** (2000), 016

[14] Brylinski, J.-L.: *Loop Spaces, Characteristic Classes and Geometric Quantization.* Prog. Math. **107**, Birkhäuser, Boston 1993

[15] Brylinski, J.-L.: *Gerbes on complex reductive Lie groups.* arXiv:math.DG/000 2158.

[16] Cardy, J. L.: *Boundary conditions, fusion rules and the Verlinde formula.* Nucl. Phys. **B 324** (1989), 581-598

[17] Carey, A. L., Johnson, S., Murray, M. K,: *Holonomy on D-Branes.* arXiv:hep-th/0204199

[18] Carter, J. S., Flath, D. E., Saito, M.: *The Classical and Quantum 6j-Symbols.* Princeton U. Press, Princeton 1995

[19] Chatterjee, D. S.: *On gerbs.* Ph.D. thesis, Trinity College, Cambridge, 1998.

[20] Di Francesco, P., Mathieu, P., Sénéchal, D.: *Conformal Field Theory.* Springer-Verlag, New York 1997

[21] Douglas, M.: *D-branes and discrete torsion.* hep-th/9807235

[22] Douglas, M.: *Topics in D-geometry.* Class. Quant. Grav. **17** (2000), 1057-1070

[23] Douglas, M.: *Lectures on D-branes on Calabi-Yau manifolds.* ICTP Lect. Notes, VII, Trieste 2002, http://www.ictp.trieste.it∼pub_off/lectures/vol7.html

[24] Douglas, M., Fiol, B.: *D-branes and discrete torsion II.* arXiv:hep-th/9903031

[25] Felder, G., Fröhlich, J., Fuchs, J., Schweigert, C.: *Conformal boundary conditions and three-dimensional topological field theory.* Phys. Rev. Lett. **84** (2000), 1659-1662

[26] Elitzur, S., Sarkissian, G.: *D-Branes on a gauged WZW model.* Nucl.Phys. **B 625** (2002), 166-178

[27] Felder, G., Gawędzki, K., Kupiainen, A.: *Spectra of Wess-Zumino-Witten models with arbitrary simple groups.* Commun. Math. Phys. **117** (1988), 127-158

[28] Fredenhagen, S., Schomerus, V.: *Branes on group manifolds, gluon condensates, and twisted K-theory.* J. High Energy Phys. **JHEP04** (2001), 007

[29] Freed, D. S., Witten, E.: *Anomalies in string theory with D-branes.* Asian J. Math. **3** (1999), 819-851

[30] Fuchs, J., Huiszoon, L. R., Schellekens, A. N., Schweigert, C., Walcher, J.: *Boundaries, crosscaps and simple currents.* Phys. Lett. **B 495** (2000), 427-434

[31] Fuchs, J., Kaste, P., Lerche, W., Lutken, C., Schweigert, C., Walcher, J: *Boundary fixed points, enhanced Gauge symmetry and singular bundles on K3.* Nucl. Phys. **B598** (2001), 57-72

[32] Fuchs, J., Runkel, I., Schweigert, C.: *TFT construction of RCFT correlators III: Simple currents.* arXiv:hep-th/0403157

[33] Fuchs, J., Schweigert, C.: *The action of outer automorphisms on bundles of chiral blocks.* Commun. Math. Phys. **206** (1999), 691-736

[34] Gaberdiel, M. R., Gannon, T.: *Boundary states for WZW models.* Nucl. Phys. **B 639** (2002), 471-501

[35] Gaberdiel, M. R., Gannon, T.: *D-brane charges on non-simply connected groups.* J. High Energy Phys. **JHEP04** (2004), 030

[36] K. Gawędzki, *Topological actions in two-dimensional quantum field theories*, In: *Non-perturbative Quantum Field Theory*, eds. 't Hooft, G., Jaffe, A., Mack, G., Mitter, P. K., Stora, R., Plenum Press, New York 1988, pp. 101-142

[37] Gawędzki, K.: *Conformal field theory: a case study.* In: *Conformal Field Theory: New Non-Perturbative Methods in String and Field Theory*, eds. Nutku, Y., Saclioglu, C., Turgut, T., Perseus 2000, pp. 1-55

[38] Gawędzki, K.: *Boundary WZW, G/H, G/G and CS theories.* Ann. Henri Poincaré **3** (2002), 847-881

[39] Gawędzki, K., Reis, N.: *WZW branes and gerbes.* Rev. Math. Phys. **14** (2002), 1281-1334

[40] Gawędzki, K., Reis, N.: *Basic gerbe over non-simply connected compact groups.* J. Geom. Phys. **50** (2004), 28-55

[41] Gawędzki, K., Todorov, I., Tran-Ngoc-Bich, P.: *Canonical quantization of the boundary Wess-Zumino-Witten model.* arXiv:hep-th/0101170, to appear in Commun. Math. Phys.

[42] Gepner, D., E. Witten, E.: *String Theory on Group Manifolds*, Nucl. Phys. **B 278** (1986), 493-549

[43] Giraud, J.: *Cohomologie non-abélienne.* Grundl. vol. bf 179, Springer 1971

[44] Hitchin, N. J.: *Lectures on special Lagrangian submanifolds.* In: *Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifold*, eds. Vafa, C., Yau, S.-T., AMS/IP Stud. Adv. Math. vol. 23, Amer. Math. Soc., Providence, RI 2001, pp. 151-182

[45] Kac, V. G.: *Infinite Dimensional Lie Algebras.* Cambridge University Press, 1985

[46] Kapustin, A.: *D-branes in a topologically nontrivial B-field.* Adv. Theor. Math. Phys. **4** (2000), 127-154

[47] Klimčik, C., Ševera, P.: *Open Strings and D-branes in WZNW model.* Nucl. Phys. **B488** (1997), 653-676

[48] Kontsevich, M.: *Mirror symmetry in dimension 3.* Séminaire Bourbaki, 801, Astérisque **237** (1996), 275-293

[49] B. Kostant: *Quantization and unitary representations.* Lecture Notes in Math., vol. 170, Springer, Berlin 1970, pp. 87-207

[50] Kreuzer, M, Schellekens, A., N.: *Simple currents versus orbifolds with discrete torsion - a complete classification.* Nucl. Phys. **B 411** (1994), 97-121

[51] Mackaay, M.: *A note on the holonomy of connections in twisted bundles*, math.DG/0106019

[52] Matsubara, K., Schomerus, V.,Smedback, M.: *Open strings in simple current orbifolds.* Nucl. Phys. B O**626** (2002), 53-72

[53] Meinrenken, E.: *The basic gerbe over a compact simple Lie group.* L'Enseignement Mathematique **49** (2003), 307-333

[54] Moore, G.: *K-Theory from a physical perspective.* arXiv:hep-th/0304018

[55] Moore, G., Seiberg, N.: *Lectures on RCFT.* Physics, Geometry, and Topology, Plenum Press, New York 1990

[56] Murray, M. K.: *Bundle gerbes.* J. London Math. Soc. (2) **54** (1996), 403-416

[57] Murray, M. K., Stevenson, D.: *Bundle gerbes: stable isomorphisms and local theory.* J. London Math. Soc. (2) **62** (2000), 925-937

[58] Pawełczyk, J.: *SU(2) WZW D-branes and their non-commutative geometry from DBI action.* J. High Energy Phys. **JHEP08** (2000), 006

[59] Petkova, V. B., Zuber, J.-B.: *Conformal boundary conditions and what they teach us.* In: *Proceedings of Nonperturbative Quantum Field Theoretic Methods and their Applications*, Horvath, Z., Palla, L. eds., World Scientific, Singapore 2001, pp. 1-35

[60] Picken, R.: *TQFT's and gerbes.* In: *Algebraic and Geometric Topology*, **4** (2004), 243-272

[61] Polchinski, J.: *TASI lectures on D-branes.* arXiv:hep-th/9611050

[62] Reis, N.: *Geometric interpretation of boundary conformal field theories.* Ph.D. thesis. ENS-Lyon 2003

[63] Runkel, I.: *Boundary structure constants for the A-series Virasoro minimal models.* Nucl. Phys. **B 549** (1999), 563-578

[64] Runkel, I.: *Structure constants for the D-series Virasoro minimal models.* Nucl. Phys. **B 579** (2000), 561-589

[65] Schellekens, A. N.: *The program Kac.* http://www.nikhef.nl∼t58/kac

[66] Schweigert, C., Fuchs, J., Walcher, J.: *Conformal field theory, boundary conditions and applications to string theory.* In: *Non-Perturbative QFT Methods and Their Applications*, Horvath, Z., Palla, L. eds., World Scientific, Singapore 2001, pp. 37-93 arXiv:hep-th/0011109

[67] Schomerus, V.: *Lectures on branes in curved backgrounds.* Class. Quant. Grav. **19** (2002), 5781-5847

[68] Sharpe, E. R.: *Discrete torsion and gerbes I, II.* arXiv:hep-th/9909108 and 9909120

[69] Vafa, C.: *Modular invariance and discrete torsion on orbifolds.* Nucl. Phys. **B 273** (1986), 592-606

[70] Walcher, J.: *Worldsheet boundaries, supersymmetry, and quantum geometry.* ETH dissertation No. 14225, 2001

[71] Witten, E.: *Non-abelian bosonization in two dimensions.* Commun. Math. Phys. **92** (1984), 455-472

[72] Witten, E.: *Overview of K-theory applied to strings.* J. Mod. Phys. **A16** (2001), 693-706